La Granja: Revista de Ciencias de la Vida
pISSN:1390-3799; eISSN:1390-8596
https://doi.org/10.17163/lgr.n44.2026.08

Review / Reseña bibliográfica

Agricultural Sciences

CC BY NC SA

Integrated Management of *Zabrotes subfasciatus* in Common Bean (*Phaseolus vulgaris*), Challenges and Control Strategies for Food Security

Manejo integrado de *Zabrotes subfasciatus* en el frijol común (*Phaseolus vulgaris*), desafíos y estrategias de control para la seguridad alimentaria

Luis David Jarquín-Izaguirre*[1], Luisa Fernanda Correa Díaz[1] and Saul Alejandro Jarquín-Izaguirre[2]

[1]*Programa de Maestría en Agricultura Tropical Sostenible, Departamento de Posgrado, Universidad Zamorano, 04001, Tegucigalpa, Honduras. [https://ror.org/01kt2cx88]*
[2]*Departamento de Agroindustria, Universidad Nacional Autónoma de Honduras, 04001, El Paraíso, Honduras. [https://ror.org/03xyve152]*

*Corresponding author: luis.jarquin@est.zamorano.edu

Article received on November 14th, 2024. Accepted, after review, on October 24th, 2025. Published on September 1st, 2026.

Abstract

Integrated management of *Zabrotes subfasciatus* in common bean (*Phaseolus vulgaris*) is crucial due to its wide distribution and the significant damage it causes to stored grains. The goal of integrated management is to reduce losses and ensure food security. This review aimed to understand the biology of *Z. subfasciatus*, identify effective control methods, and propose an integrated model that minimizes the use of chemical insecticides. A systematic review of scientific literature published since 2000 was conducted, evaluating 653 initial studies, of which 61 met the inclusion criteria. Key aspects such as taxonomy, geographic distribution, life cycle, and pest control methods were addressed using information from databases such as ScienceDirect, Elsevier, and SciELO. The main findings showed that an integrated management approach combining cultural, physical, chemical, and genetic methods is effective in controlling the pest without relying solely on chemical insecticides, whose indiscriminate use poses risks to human health and the environment. Monitoring and the development of resistant bean varieties stand out as fundamental practices in pest management. In conclusion, integrated management of *Z. subfasciatus* significantly contributes to the protection of stored grains and, consequently, to the food security of farming communities. It is recommended to strengthen farmer training in sustainable control techniques and to promote research on alternative management methods.

Keywords: Storage, pest control, entomology, bean, insect.

©2026, Universidad Politécnica Salesiana, Ecuador.

Resumen

El manejo integrado del *Zabrotes subfasciatus* en frijol común (*Phaseolus vulgaris*) es crucial, debido a su amplia distribución y el daño significativo que causa en granos almacenados. Su manejo integrado busca reducir las pérdidas y garantizar la seguridad alimentaria. El objetivo de esta revisión es comprender la biología de *Z. subfasciatus*, identificar métodos de control eficaces y proponer un modelo integral que reduzca el uso de insecticidas químicos. Se realizó una revisión sistemática de literatura científica publicada a partir del año 2000, evaluando 653 estudios iniciales, de los cuales 61 cumplieron con los criterios de inclusión. Se abordaron aspectos como la taxonomía, distribución geográfica, ciclo de vida y métodos de control de la plaga, a partir de información recopilada en bases de datos como ScienceDirect, Elsevier y SciELO. Los resultados principales evidenciaron que un enfoque de manejo integrado, que combine métodos culturales, físicos, químicos y fitogenéticos, es eficaz para controlar la plaga sin depender exclusivamente de insecticidas químicos, cuyo uso indiscriminado representa un riesgo para la salud humana y el ambiente. El monitoreo y el desarrollo de variedades resistentes de frijol se destacan como prácticas fundamentales en el manejo de la plaga. En conclusión, el manejo integrado de *Z. subfasciatus* aporta significativamente a la protección de los granos almacenados y, por ende, a la seguridad alimentaria de las comunidades agrícolas. Se recomienda fortalecer la capacitación de los agricultores en técnicas sostenibles de control y promover la investigación en métodos alternativos de manejo.

Palabras clave: Almacenamiento, control de plaga, entomología, frijol, insecto.

Suggested citation: Jarquín, L., Correa, L. and Jarquín, S. (2026). Integrated Management of *Zabrotes subfasciatus* in Common Bean (*Phaseolus vulgaris*), Challenges and Control Strategies for Food Security. La Granja: Revista de Ciencias de la Vida. Vol. 44(2):136-150. https://doi.org/10.17163/lgr.n44.2026.08.

Orcid IDs:

Luis David Jarquín-Izaguirre: https://orcid.org/0009-0001-9970-8548
Luisa Fernanda Correa Díaz: https://orcid.org/0009-0001-3577-2399
Saul Alejandro Jarquín-Izaguirre: https://orcid.org/0009-0002-3819-099X

©2026, Universidad Politécnica Salesiana, Ecuador.

1 Introduction

The common bean (*Phaseolus vulgaris*) is the most important legume used for direct human consumption (Didinger et al., 2022). A member of the Fabaceae family, the common bean is a predominantly self-pollinating species native to Mesoamerica (Bitocchi et al., 2017). Due to its wide geographic distribution, this ancient legume split and evolved into two domesticated gene pools: the Mesoamerican and the Andean. The highest-quality varieties were developed and domesticated in South America, based on the Andean gene pool, while the lower-quality varieties originated in Mexico and Central America, with roots in the Mesoamerican gene pool (Wilker et al., 2020). Thanks to this genetic diversity and its adaptability, the common bean has established itself as one of the most important crops for human consumption worldwide (Refaay et al., 2021). Its high nutritional value makes it an essential food, especially in Latin America and developing countries (Hernández-Ochandía et al., 2018), where it constitutes an important source of protein, vitamins, and minerals. Dried beans contain approximately 22 % protein, complex carbohydrates (62 %), soluble fiber (15 %), and various micronutrients such as calcium, iron, magnesium, phosphorus, potassium, and folate, positioning them as a key food in balanced and healthy diets (Myers and Kmiecik, 2017).

In terms of production, *P. vulgaris* accounts for 85 % of global bean production, making it the most widely cultivated legume on the planet (Mamo et al., 2023). Annually, more than 27 million tons of beans are harvested on approximately 29 million hectares worldwide, feeding more than 300 million people who depend on agricultural economies in various regions of the world (Nasar et al., 2023). Latin America stands out as the region with the highest production and consumption, accounting for 45 % of global production (IICA / Proyecto Red SICTA, 2013). In this region, beans are not only a staple food but also a vital crop for the economy of smallholder farmers. Particularly in Central America, beans occupy a strategic position in food security and the rural economy. According to FAOSTAT data, in 2022 Honduras produced 130 906 tons of beans, with an average yield of 802 kg/ha, which highlights their importance within the region's agricultural economies (FAO, 2022).

Despite its importance of economic and food security, the crop faces various biotic and abiotic stresses that limit its productivity. Among biotic factors, the storage pests *Zabrotes subfasciatus* and *Acanthoscelides obtectus* stand out as the most destructive during the postharvest stage (Tigist et al., 2021). In particular, *Z. subfasciatus* represents one of the main threats to producers, causing significant losses that affect both commercialization and the availability of seeds for future plantings (Mukankusi et al., 2019). The management of these pests in stored grains has traditionally relied on the application of synthetic insecticides such as phosphine, pyrethroids, and organophosphate compounds. However, many of these products have limited availability to farmers in developing countries (Agrafioti et al., 2019; Gourgouta et al., 2019). Furthermore, the continued and indiscriminate use of pesticides has generated growing global concern due to the risks it poses to human health and the environment, highlighting the urgency of implementing more sustainable strategies for pest management in stored products (Souza et al., 2010).

Given its fundamental role in food security, the rural economy, and the nutrition of millions of people in developing countries, the common bean faces the challenge of maintaining its productivity in the face of threats such as *Zabrotes subfasciatus*, one of the main postharvest pests. Ensuring its sustainable management is key to reducing losses, preserving grain quality, and strengthening the livelihoods of smallholder farmers. In this context, this study aims to understand the biology of *Z. subfasciatus*, identify effective control methods, and propose an integrated management model that minimizes the use of chemical insecticides, thereby contributing to crop protection and the sustainability of agricultural systems.

2 Materials and Methods

2.1 Selection of Studies

Initially, a selection of inclusion criteria was carried out, identifying topics of interest within the field of bean pest control, with an emphasis on *Zabrotes subfasciatus*. Based on this identification, the key parameters for the literature search were defined. The inclusion criteria considered were: a) articles addressing the taxonomy of *Z. subfasciatus*, b) studies

©2026, Universidad Politécnica Salesiana, Ecuador.

on its biology and life cycle, c) research related to its geographic distribution, and d) control methods (cultural, phytogenetic, physical-mechanical, and chemical). In general, priority was given to literature published since 2000; however, for specific topics such as taxonomy, morphology, and physiology, earlier studies were included if they were relevant in terms of content and scientific validity.

2.2 Identification of Databases

The repositories used to identify scientific articles were ScienceDirect, Elsevier, and SciELO. Once an article met the inclusion criteria, the literature was analyzed and synthesized, involving a critical evaluation, the extraction of relevant data and scientific evidence, and the identification of trends and fundamental information about the species.

2.3 Analysis Structure

A basic structure was defined to organize the collected information, covering aspects such as taxonomy, morphology, geographic distribution, biology and life cycle, monitoring, as well as control methods in storage: cultural, phytogenetic, physical-mechanical, and chemical. This organization allowed for a comprehensive approach to the management of *Zabrotes subfasciatus* in *Phaseolus vulgaris*.

2.4 Final Selection of Studies for the Analysis

For the selection of articles, the PRISMA approach was used to guide the process of identifying, selecting, and documenting the studies included in this systematic review, following the methodology framework of Page et al. (2021), initially identifying 653 records through searches in scientific databases. Subsequently, the methodological quality of these studies was assessed using a checklist. This tool allowed us to identify potential biases and evaluate the internal validity, methodological clarity, and relevance of the studies. After removing duplicate articles, those that did not meet the thematic or methodological criteria were excluded, resulting in a total of 61 studies included for analysis (Figure 1).




Figure 1. PRISMA flow diagram for the systematic review of *Zabrotes subfasciatus* in beans.

©2026, Universidad Politécnica Salesiana, Ecuador.

3 Results and Discussion

3.1 General characteristics of bean cultivation

The genus *Phaseolus* spp. consists of 76 species, 5 of which are domesticated species, including *P. vulgaris* (Kaplan, 2003). This genus is notable for its cultivated species, which occupy 90 % of the land dedicated to bean cultivation worldwide (Singh, 2013). It is a herbaceous plant that was independently domesticated in ancient Mesoamerica and the Andes (Kaplan, 1981). It is classified within the group of dicotyledonous plants (Katungi et al., 2009). The seeds can be consumed immature (tender), mature, fresh, or dried (Morales-Santos et al., 2017).

There are two bean planting cycles per year, but farmers mainly choose to plant during the late season (October–December). Planting during the early season (May–September) is less common, as it leads to grain losses due to germination or rot (Torres-Treminio and Mendoza-Montoya, 2002). Post-harvest handling of common beans is a critical stage in the agricultural production chain, where beans are exposed to various factors that can compromise their quality and quantity. In Central America and Mexico, losses of up to 35 % of stored beans are estimated (Espinal et al., 2021).

3.2 Taxonomy of *Zabrotes subfasciatus*

The Mexican bean weevil or bean beetle, *Zabrotes subfasciatus*, belongs to the order Coleoptera, a species of leaf beetle in the family Chrysomelidae. It is the primary phytosanitary problem affecting stored beans in areas below 1 500 meters in altitude (Cárdenas-Morales et al., 2019). It is characterized by complete metamorphosis and feeds exclusively on seeds (Dobie et al., 1991).

Members of this family are easily recognizable; the body, covered with small hairs, is compact and globular, the elytra are short, and the pygidium is exposed (Rees, 2018). Their primary host is *Phaseolus vulgaris* in its seed stage; however, they can also feed on other legume species under isolated conditions, such as *Vigna unguiculata* L. (Ramírez Cariño et al., 2017).

Table 1. Taxonomy of *Zabrotes subfasciatus*.

Taxonomic classification	
Taxonomic category	**Taxon to which it belongs**
Domain	Eukarya
Kingdom	Animalia
Phylum	Arthropoda
Class	Insecta
Order	Coleoptera
Suborder	Polyphaga
Superfamily	Chrysomeloidea
Family	Chrysomelidae
Genus	*Zabrotes*
Species	*Z. subfasciatus* (Boheman, 1833)
Common name	Mexican or pinto weevil

Source: Population increase of *Zabrotes subfasciatus* in common bean cultivars (Cárdenas-Morales et al., 2019).

©2026, Universidad Politécnica Salesiana, Ecuador.

3.3 Morphology of *Zabrotes subfasciatus*

This beetle exhibits distinctive characteristics in both males and females, which are crucial for its identification and study. The adult body is oval-shaped, thick, convex, and gray-brown or black in color, except for the base of the antennae and the tips of the tarsi. The body measures approximately 2 to 3 mm in length and 1 to 2 mm in width. Its antennae are long and can reach 1.45 mm in length; the segments are long and black, except for the first two, which are reddish (Dell'Orto and Arias-Velásquez, 1985). The elytra are short and relatively wide, square in shape, with the hind wings concealed (Dobie et al., 1991).

The female is larger than the male and is characterized by four white spots on the elytra (Figure 2) (Arnal and Ramos, 2006). The male is a uniform gray-brown color (Figure 3). Its hind femur is smooth, without protuberances or teeth. The part of the tibia near the femur has two movable spurs (Dobie et al., 1991). The larvae are robust, curved, and ivory-white. They are very voracious, and several may be found in a single grain (Figure 4) (Carrera-Rodríguez et al., 2009).



Figure 2. Dorsal and lateral view of the female *Zabrotes subfasciatus* (Johnson, 2019).



Figure 3. Dorsal and lateral views of the male *Zabrotes subfasciatus* (Johnson, 2019).

©2026, Universidad Politécnica Salesiana, Ecuador.



Figure 4. Larva of *Zabrotes subfasciatus* (Johnson, 2019).

3.4 Morphology of the Male

A beetle with a mostly black body, with some lighter areas in certain individuals. The head has white pubescence with a yellowish-brown spot on the *vertex*. The pronotum and elytra have small, scattered brown and yellow spots. The abdomen is mostly white, with brown patches on the sides. The head structure shows fine punctures, with a semicircular pronotum and uniformly punctured elytra. Measurements vary, but the beetle generally has a length of 1.56 to 2.16 mm and a width of 1.26 to 1.62 mm. Regarding the genitalia, the median lobe is wider at the base and has a subtriangular ventral valve. The lateral lobes are shorter and feature an apical cleft (Nápoles, 2018).

3.5 Morphology of the Female

Sexual dimorphism is present in this species, with females having a median band on the pronotum and a small white spot in each corner, as well as a transverse white band on the elytra. However, pubescence is generally similar to that of males. Antennal length is slightly shorter, ranging from 0.6 to 0.63 times body length. Typical measurements of length (pronotum-elytra) range from 2.28 to 2.46 mm, with a width of 1.74 to 1.86 mm and a maximum thoracic depth of 1.26 to 1.32 mm (Nápoles, 2018).

3.6 Geographic Distribution of *Zabrotes subfasciatus*

This species' ability to thrive under storage conditions has made it widely available worldwide (Del Obregón and Gómez-Gutiérrez, 2023). This insect is widely distributed in the tropical and subtropical regions of Africa, Madagascar, India, and the Americas, where it is believed to be native (Dobie et al., 1991). In the Americas, it is distributed from the United States to Brazil (Arnal and Ramos, 2006). It is mainly found in warm, humid climates at low elevations above sea level (Dell'Orto and Arias-Velásquez, 1985).

3.7 Reproductive Habits of *Zabrotes subfasciatus*

The female mates with the male, and this mating appears to be influenced by bean seeds. Upon contact with these, the number of sexual encounters increases, stimulating oogenesis (egg development) and oviposition activity. It has been documented that olfactory, gustatory, and tactile chemoreceptors regulate or control the reproductive activity of females (Pimbert and Pierre, 1983).

Females oviposite on the naked seed, never on the pods (Credland and Dendy, 1992). However, Pimbert and Pierre (1983) note that they are capable of reproducing on both parts in the field and in storage. When the pods of *P. vulgaris* and *P. lunatus* are partially or fully dehiscent, this allows the female to encounter the seeds to lay eggs. These same

©2026, Universidad Politécnica Salesiana, Ecuador.

authors note that *Z. subfasciatus* cannot pierce the pods to lay eggs on the seeds, nor can it lay eggs inside the pods as other members of the family can. If the female deposits the eggs on the walls of the pods, they adhere, and the larva can penetrate the walls but dies inside the pod since it cannot penetrate the seed (Del Obregón and Gómez-Gutiérrez, 2023).

3.8 Life Cycle of *Zabrotes subfasciatus*

3.8.1 Eggs

Females lay between 35 and 60 eggs and infest up to 36 grains during their reproductive cycle; the incubation period can range from four to eight days, followed by the larval stage (12 days). The eggs measure approximately 0.3 mm in diameter and are rounded. Two days after emerging, the female *Z. subfasciatus* usually begins to lay eggs (Credland and Dendy, 1992). Studies by Golob and Kilminster (1982) indicate that the female has a 1-day preoviposition period; subsequently, the female lays 50 eggs during her 13-day lifespan, with the majority laid after the fourth day. The egg is strongly attached to the grain's testa by a viscous, brightly colored, translucent substance that acts as glue. This glue or secretion that attaches the eggs is generally deposited a few seconds before oviposition (Credland and Dendy, 1992).

3.8.2 Larvae

They are vermiform and apodal, measuring 3 to 3.5 mm in length. Their body tends to be curved, with several folds, whitish in color, and a reduced head embedded in the prothorax (Ascencio-Alvarado, 2012). Newly emerged larvae penetrate the interior of the grain after consuming most of the cotyledon, constructing a tunnel and leaving only a covering over the grain's pericarp through which the adult subsequently emerges (Dell'Orto and Arias-Velásquez, 1985).

3.8.3 Pupae

They measure about 3 mm in length. They are of the exarate type, with appendages. The difference between the sexes can be noted at this stage by the shape of the last abdominal segment, which is straight in the female and arched in the male (Bonet et al., 2005; Ascencio-Alvarado, 2012). This stage lasts between four and nine days, after which the larva enters the adult stage and subsequently emerges through a hole it makes in the testa of the grain (González et al., 1984; Bonet et al., 2005).

3.8.4 Adults

Females, which are larger than males, can be easily distinguished by the presence of four light spots on the pronotum, which contrast with the dark, shiny color of the body. They have well-developed wings, and their body is oval-shaped with a free-standing head. The eyes are well-developed, and their antennae have 11 segments. The hind legs are thicker than the front ones (Bonet et al., 2005; Ascencio-Alvarado, 2012).

The exit point of the adult resembles a circular window on the seed surface (Rees, 2018). Emerging adults do not need to feed before or after mating, as they acquire all the necessary nutrients to complete their life cycle during their larval development. They are highly active, capable of running and flying quickly, and are frequently observed moving across the surface of infested seeds.



Figure 5. Life cycle of *Zabrotes subfasciatus* (CABI, 2019).

©2026, Universidad Politécnica Salesiana, Ecuador.

3.9 Infection Conditions

Its life cycle lasts 24 to 25 days under conditions of 32 °C and 70 % relative humidity (Dobie et al., 1991). However, other authors indicate that it takes more than 30 days from egg to adult (Golob and Kilminster, 1982; Credland and Dendy, 1992; Barbosa et al., 1999). The optimal temperature for development is 37–38 °C, with a minimum slightly below 20 °C (Dobie et al., 1991). Adults have a short lifespan. Various authors have determined the average lifespan of adults. For example, Dell'Orto and Arias-Velásquez (1985) indicated that the adult *Z. subfasciatus* lives for about 10 to 12 days, while Barbosa et al. (1999) argue that it lives for about 36 days.

3.10 Feeding and Damage

It is the most significant pest of stored common beans and lima beans (*P. lunatus*) (Dobie et al., 1991). It can also feed on other legume seeds, such as cowpeas, peas, lentils, and soybeans (Dell'Orto and Arias-Velásquez, 1985; Rees, 2018). The damage caused by this insect species results from the larvae invading and feeding inside the grains, leading to weight loss, reduced nutritional value, and compromised product hygiene due to the presence of excrement, eggs, and insects. Additionally, the germination capacity of the seeds may be reduced or completely lost (Barbosa et al., 1999).

3.11 Cultural Control

A traditional method used to control the population of *Z. subfasciatus* is to store bean seeds inside their own pods (Arnal and Ramos, 2006; Del Obregón and Gómez-Gutiérrez, 2023). This is because females can generally only lay eggs on bare seeds and very rarely on pods unless they are dehiscent (Pimbert and Pierre, 1983).

In many places, a mixture of local plants is used with stored grain. Information about which plants should be used for pest control and which parts of them should be mixed with the grain is passed down from generation to generation in rural areas (Carrera-Rodríguez et al., 2009). Plant-derived substances have been used for pest control since ancient times. However, their use declined with the advent of chemical insecticides, becoming limited to rural areas (Rodríguez Hernández and López Pérez, 2001).

In developing countries, it is common to mix plant-based products with stored beans to protect them during storage. Among the products that have been used are neem leaves (*Azadirachta indica*), *Ocimum canum* leaves, *Capsicum frutescens* fruits, and *Piper nigrum* fruits (Madueke-Onu et al., 2015; Perales-Aguilar et al., 2020; Okafor et al., 2024). Another method involves the use of vegetable oils, generally of local origin. Among the vegetable oils that have been found to be effective are those derived from the seeds of *Anthonomus grandis* (Perales-Aguilar et al., 2020). However, a disadvantage of using oils is that they impart a rancid and sticky appearance to the grains, which could reduce market acceptance (Álvarez Serrano et al., 2021).

The use of mineral powders as a cultural control method for stored grain insects, including *Z. subfasciatus*, has been widely employed (Lopez-Monzon et al., 2016). Mixing sand or ash with the grains is an alternative method for insect control. Sand or ash damages the insect's cuticle by scraping it, causing it to lose moisture. If the grain is dry, there will not be enough moisture for the insect to replenish what it has lost, so it will die from desiccation (Kalpna et al., 2022); ash creates a physical barrier for weevils.

Mixing lime with stored grains is a common practice in northwestern Mexico. Permuy-Abeleira et al. (2008) report that farmers in the region clean the storage area before harvests, removing any debris found, and wash the area with soap; after the grain is harvested, it is mixed with lime to prevent insect attacks.

The use of garlic (*Allium sativum*) extracts as a natural repellent and insecticide against *Zabrotes subfasciatus* in stored beans has also been documented. Studies conducted in El Salvador have shown that aqueous and hydroalcoholic garlic extracts, applied at concentrations between 5 % and 20 %, achieve significant effects on both adult mortality and infestation reduction, without compromising grain quality (Rámos-Archila and Santacruz-Jiménez, 2024).

Another method used is exposing the grains to the sun. The insects abandon the grain when it is exposed to sunlight due to the high temperatures (above 40–44 °C). However, sun exposure does not always kill the eggs and larvae that remain insi-

©2026, Universidad Politécnica Salesiana, Ecuador.

de the grain (Carrera-Rodríguez et al., 2009). This method only protects the outer layers of the grain, so it can only harm adult insects on the surface or larvae that are moving freely (Cuevas, 2006).

3.12 Phytogenetic Control

A wide variety of proteins found in the seeds of wild *P. vulgaris* plants (arcelin and α-amylase inhibitors) have been used to control *Z. subfasciatus* in stored beans (Goossens et al., 2000). Although bean seeds possess defense proteins such as enzyme inhibitors and secondary metabolites (alkaloids, saponins), they can still be attacked. This is because insects co-evolve with their food sources, and in the course of evolution, new insect strains emerge that allow them to defend themselves against the seeds' natural chemicals (Rodríguez-Sifuentes et al., 2020).

3.13 Physical-Mechanical Control

In the tropical regions of some Central American countries, as well as in Mexico, airtight storage of grains for human consumption is commonly used; under these conditions, insects experience stress due to reduced oxygen levels, creating an unfavorable environment for their survival (Moreno-Martinez et al., 2000). Low levels of O_2 and CO_2 cause metabolic stress in insects, generally increasing mortality and decreasing fecundity and development rate (Thompson and Suarez, 2009).

Another physical-mechanical treatment involves heat, which consists of heating a container to a specific temperature (50 to 55 °C) for 20 or 30 hours. This duration allows the container materials to expand and adjust to the change. Although storage insects die in less than an hour at these temperatures, such a long duration is necessary because the heat must penetrate to where the insects might be found. However, these control methods are not typically cost-effective, as the containers or heating equipment are often expensive.

3.14 Chemical Control

This is the most widely used method in grain warehouses; among the most commonly used are fumigantes (Perales-Aguilar et al., 2020). Phosphamine fumigants and methyl bromide are widely used in the control of stored grain insects (Dierksmeier-Corcuera, 2007). In highly technical warehouses, it is common to use insecticides to control pests that affect stored products. This approach involves the application of chemicals, which can take different forms, such as powder, liquid, or fumigants.

Powdered products are particularly used in warehouses where grain is stored in bulk or in bags, and their application can be both preventive and corrective. The most popular insecticides in this category include malathion, pirimiphos-methyl, and dichlorvos, among others. Although some of these products can also be used as sprays, the use of pyrethroid insecticides is widely preferred, with deltamethrin being particularly prominent. Additionally, many warehouses rely on fumigants, with aluminum phosphide being one of the most prominent (García-Lara et al., 2007).

Vázquez-Zamorano (2014) conducted a study to evaluate the effectiveness of permethrin, diazinon, and cypermethrin in controlling adult weevils. Permethrin was found to be the most effective insecticide, with a mortality rate exceeding 60 % when using the highest dose of 500 ppm. In a study conducted in Argentina to evaluate the efficacy of aluminum phosphide in the bag-in-silo storage system, it was observed that although there is some loss of the fumigant, by hermetically sealing the storage system, it is capable of retaining the adequate amount of fumigant to eliminate stored-product insects (Carpaneto et al., 2016).

The use of chemical control methods poses significant problems: risk of toxic residues in food, insect resistance, negative environmental impact, and increasing regulatory restrictions, in addition to the costs associated with their purchase, application, and specialized equipment (Cevallos-Taxi, 2020). From an economic standpoint, chemical control may be more effective in scenarios involving large volumes and mechanized storage facilities, but it is less accessible and viable in rural areas with limited resources (Raygoza-Martínez et al., 2025).

In contrast, cultural control has proven to be highly cost-effective in rural communities, especially when based on traditional knowledge and local resources. This effectiveness stems from the fact that it utilizes locally available inputs, requires

©2026, Universidad Politécnica Salesiana, Ecuador.

little initial investment, does not depend on external commercial products, and can be easily adopted by farming families, thereby reducing risks to human health and the environment.

4 Conclusions

The wide geographic distribution of *Zabrotes subfasciatus* and its ability to infest various legumes highlight the need for ongoing research and the implementation of monitoring and control practices adapted to different environmental and cultural contexts. The integrated management approach, which combines cultural, physical-mechanical, chemical, and phytogenetic methods, represents an effective and sustainable strategy for controlling this pest, minimizing dependence on chemical insecticides and their associated risks.

However, challenges remain, such as the limited adoption of sustainable practices, the lack of accessible technologies, and inadequate storage infrastructure. For this reason, it is recommended to deepen research into the development of resistant varieties as a key measure to ensure protein availability and contribute to food security. Similarly, efforts should be made to promote the development of new biological control alternatives and innovation in postharvest management technologies, incorporating participatory approaches that address the needs of smallholder farmers.

It is also crucial that public agricultural policies integrate post-harvest management and strengthen technical assistance programs with a focus on promoting sustainable practices as part of rural development and food security strategies. This involves promoting training programs, incentives for agroecological practices, and improvements in extension systems.

Declaration on the Use of Artificial Intelligence

The authors DECLARE that, during the preparation of the article entitled "Integrated Management of *Zabrotes subfasciatus* in Common Bean (*Phaseolus vulgaris*), Challenges and Control Strategies for Food Security", no generative artificial intelligence tools or automated assistance systems were used for the writing, analysis, data interpretation, content generation, translation, or editing of the manuscript.

The authors assume full responsibility for the content, originality, integrity, and final version of the article.

Author Contributions

L.D.J.I.: Conceptualization, Data curation, Investigation, Supervision, Writing – review and editing. **L.F.C.D.:** Investigation, Data curation, Methodology, Writing – original draft. **S.A.J.I.:** Conceptualization, Formal analysis, Validation, Writing – original draft.

References

Agrafioti, P., Athanassiou, C. G., and Nayak, M. K. (2019). Detection of phosphine resistance in major stored-product insects in greece and evaluation of a field resistance test kit. *Journal of Stored Products Research*, *82*, 40–47. Online: https://doi.org/10.1016/j.jspr.2019.02.004

Álvarez Serrano, A., Milan-Leyva, E., and Matos-Castillo, Y. (2021). Estudio del efecto de aceites vegetales en el control del gorgojo (*Callosobruchus maculatus* f) en la conservación de granos de frijol caupí. *Opuntia Brava*, *13*(3), 77–86. Online: https://shortlink.uk/1wIsu

Arnal, E. A., and Ramos, F. A. (2006). *Guía práctica sobre alternativas de control de insectos-plaga en los cultivos de caraota y frijol* (Inf. Téc. n.º 12). Publicación Especial (INIA). Online: https://shortlink.uk/1rhwg

Ascencio-Alvarado, J. I. (2012). *Evaluación de la actividad insecticida y repelente del extracto obtenido a partir de la semilla de Annona diversifolia (anona) sobre el Zabrotes subfasciatus (gorgojo común del frijol)* (Bachelor's thesis, Universidad de El Salvador). Online: https://shortlink.uk/1wIsx

Barbosa, F. R., Yokoyama, M., Arraes-Pereira, P. A., and Pfeilsticker-Zimmermann, F. J. (1999). Efeito da proteína arcelina na biologia de *Zabrotes subfasciatus* (Boheman 1833), em feijoeiro. *Pesquisa*

©2026, Universidad Politécnica Salesiana, Ecuador.

Agropecuária Brasileira, *34*(10), 1805–1810. Online: https://shortlink.uk/1wIsB2

Bitocchi, E., Rau, D., Bellucci, E., Rodriguez, M., Murgia, M. L., Gioia, T., … Papa, R. (2017). Beans (*Phaseolus* ssp.) as a model for understanding crop evolution. *Frontiers in Plant Science*, *8*, 722. Online: https://doi.org/10.3389/fpls.2017.00722

Bonet, A., Morales, C. O., and Rojas, C. V. (2005). El control biológico con parasitoides: una alternativa para limitar a los gorgojos en frijol almacenado. *Fitosanidad*, *9*(2), 35–42.

CABI. (2019). *Zabrotes subfasciatus (Mexican bean weevil). CABI Compendium.* Online Publication. Online: https://doi.org/10.1079/cabicompendium.57289

Cárdenas-Morales, M., Lavastida-Pérez, M., Valdés-Herrera, R., Pozo-Velazquez, E., and Rodríguez-Rojas, R. (2019). Incremento poblacional de *Zabrotes subfasciatus* en cultivares de frijol común. *Centro Agrícola*, *46*(2), 38–45. Online: https://shortlink.uk/1wIsH

Carpaneto, B., Bartosik, R., Cardoso, L., and Manetti, P. (2016). Pest control treatments with phosphine and controlled atmospheres in silo bags with different airtightness conditions. *Journal of Stored Products Research*, *69*, 143–151. Online: https://doi.org/10.1016/j.jspr.2016.07.007

Carrera-Rodríguez, M., Martínez-González, G. M., Navarrete-Bolaños, J. L., Botello-Alverez, J. E., Rico-Martínez, R., and Jiménez-Islas, H. (2009). Estudio numérico del efecto de la temperatura ambiental en la convección natural bidimensional en el almacenamiento de granos en silos cilíndricos. *Revista Mexicana de Ingeniería Química*, *8*(1), 77–91. Online: https://shortlink.uk/1wIsM

Cevallos-Taxi, S. F. (2020). *Evaluación del efecto de ozono (o_3) en el control de gorgojo Pagiocerus frontalis (f.), en granos almacenados de maíz suave (Zea mays l.) en la provincia Cotopaxi, campus experimental Salache, proyecto granos andinos 2020* (Bachelor's thesis, Universidad Técnica de Cotopaxi). Online: https://shortlink.uk/1wIsP

Credland, P. F., and Dendy, J. (1992). Comparison of seed consumption and the practical use of insect weight in determining effects of host seed on the Mexican bean weevil, *Zabrotes subfasciatus* (Boh.). *Journal of Stored Products Research*, *28*(4), 225–234. Online: https://doi.org/10.1016/0022-474x(92)90001-7

Cuevas, M. I. (2006). Insecticidas alternativos para el control de plagas en granos almacenados. *Inventio*, *2*(3), 55–60. Online: https://shortlink.uk/1wIsV

Dell'Orto, T. H., and Arias-Velásquez, C. J. (1985). *Insectos que dañan granos y productos almacenados* (Inf. Téc. n.º 4). Serie Tecnología Postcosecha (FAO/RLAC). Online: https://shortlink.uk/1wIsZ

Del Obregón, N. C., and Gómez-Gutiérrez, O. J. (2023). Tecnologías y prácticas que utilizan los productores en el manejo poscosecha de maíz y frijol en Estelí, Madriz y Nueva Segovia, Nicaragua. *La Calera*, *23*(41). Online: https://doi.org/10.5377/calera.v23i41.16739

Didinger, C., Foster, M. T., Bunning, M., and Thompson, H. J. (2022). Nutrition and human health benefits of dry beans and other pulses. En M. Siddiq and M. A. Uebersax (Eds.), *Dry beans and pulses* (pp. 481–504). Wiley. Online: https://doi.org/10.1002/9781119776802.ch19

Dierksmeier-Corcuera, G. (2007). Características y mecanismos de acción de algunos compuestos usados en el combate de plagas de almacén. *Fitosanidad*, *11*(2), 81–84.

Dobie, P., Haines, C. P., Prevett, P. F., and Rees, D. P. (1991). *Insects and arachnids of tropical stored products: their biology and identification*. Natural Resources Institute.

Espinal, J. R., Soto, O. J., Abrego, A. N., Pizarro, I. P., Rodriguez, I. Y., and Rosas-Sotomayor, J. C. (2021). Resistencia de genotipos de frijol común a los daños causados por el gorgojo mexicano *Zabrotes subfasciatus* (Boheman) en el almacenamiento. *CEIBA (Edición Zamorano Investiga)*, 112–121. Online: https://shortlink.uk/1wIt6

FAO. (2022). *FAOSTAT database: Dry Bean Yield.* Online: https://www.fao.org/faostat/en/#data/QCL

García-Lara, S., Espinosa-Carrillo, C., and Bergvinson, D. (2007). *Manual de plagas en granos almacenados y tecnologías alternas para su manejo y control*. México, D.F.: CIMMYT.

©2026, Universidad Politécnica Salesiana, Ecuador.

Golob, P., and Kilminster, A. (1982). The biology and control of *Zabrotes subfasciatus* (Boheman) (Coleoptera: Bruchidae) infesting red kidney beans. *Journal of Stored Products Research*, *18*(3), 95–101. Online: https://doi.org/10.1016/0022-474x(82)90008-x

González, M., Roche, R., and Simanca, M. E. (1984). Ciclo de vida de *Zabrotes subfasciatus* (Coleoptera: Bruchidae), plaga de granos almacenados. *Ciencias de la Agricultura*(21). Online: https://shortlink.uk/1wItb

Goossens, A., Quintero, C., Dillen, W., Rycke, R. d., Flower Valor, J., Clercq, J. d., … Angenon, G. (2000). Analysis of bruchid resistance in the wild common bean accession G02771: no evidence for insecticidal activity of arcelin 5. *Journal of Experimental Botany*, *51*(348), 1229–1236. Online: https://doi.org/10.1093/jexbot/51.348.1229

Gourgouta, M., Rumbos, C. I., and Athanassiou, C. G. (2019). Residual toxicity of a commercial cypermethrin formulation on grains against four major storage beetles. *Journal of Stored Products Research*, *83*, 103–109. Online: https://doi.org/10.1016/j.jspr.2019.05.001

Hernández-Ochandía, D., Rodríguez-Hernández, M. G., Miranda-Cabrera, I., Moreno-León, E., Castro-Lizazo, I., Delgado-Oramas, B. P., and Holgado, R. (2018). Reproducción y efecto nocivo de *Meloidogyne incognita* (Kofoid y White) Chitwood en *Phaseolus vulgaris* L. 'cuba-cueto-25-9'. *Revista de Protección Vegetal*, *33*(2). Online: https://shortlink.uk/1wIth

IICA / Proyecto Red SICTA. (2013). *Estudio de las cadenas de valor maíz blanco y frijol en Centroamérica* (Inf. Téc. n.º 115). Instituto Interamericano de Cooperación para la Agricultura. Online: https://shortlink.uk/1wItn

Johnson, A. (2019). *Zabrotes subfasciatus (Mexican bean weevil).* CABI Compendium. Online: https://doi.org/10.1079/cabicompendium.57289

Kalpna, Hajam, Y. A., and Kumar, R. (2022). Management of stored grain pest with special reference to *Callosobruchus maculatus*, a major pest of cowpea: A review. *Heliyon*, *8*(1), e08703. Online: https://doi.org/10.1016/j.heliyon.2021.e08703

Kaplan, L. (1981). What is the origin of the common bean? *Economic Botany*, *35*(2), 240–254. Online: https://doi.org/10.1007/bf02858692

Kaplan, L. (2003). Taxonomy, Distribution, and Ecology of the Genus *Phaseolus* (Leguminosae–Papilionoideae) in North America, Mexico, and Central America. *Economic Botany*, *57*(3), 421. Online: https://shortlink.uk/1wItr

Katungi, E., Farrow, A., Chianu, J., and Beebe, S. (2009). *Common bean in Eastern and Southern Africa: A situation and outlook analysis* (Inf. Téc.). CIAT. Online: https://shortlink.uk/1wItw

Lopez-Monzon, C. E., Tobar-Tomas, W. V., and Ventura-Gomez, A. G. (2016). Controles alternativos para el gorgojo del frijol (*Acanthoscelides obtectus* Say) en granos almacenados a partir de materiales naturales y minerales. *Ciencia, Tecnología y Salud*, *3*(2), 149–156. Online: https://doi.org/10.36829/63CTS.v3i2.196

Madueke-Onu, F., Ogu, E., and Ikehi, M. (2015). Use of neem and garlic dried plant powders for controlling. *Egyptian Journal of Biological Pest Control*, *25*(2), 507–512.

Mamo, T., Singh, A., Singh, A., Mahama, A. A., and Suza, W. (2023). Common bean breeding. En *Crop improvement*. Iowa State University Digital Press. Online: https://shortlink.uk/1rhxk

Morales-Santos, M. E., Peña Valdivia, C. B., García-Esteva, A., Aguilar-Benítez, G., and Kohashi-Shibata, J. (2017). Características físicas y de germinación en semillas y plántulas de frijol (*Phaseolus vulgaris* L.) silvestre, domesticado y su progenie. *Agrociencia*, *51*(1), 43–62. Online: https://shortlink.uk/1rhxn

Moreno-Martinez, E., Jiménez, S., and Vázquez, M. E. (2000). Effect of *Sitophilus zeamays* and *Aspergillus chevalieri* on the oxygen level in maize stored hermetically. *Journal of Stored Products Research*, *36*(1), 25–36. Online: https://doi.org/10.1016/s0022-474x(99)00023-5

Mukankusi, C., Raatz, B., Nkalubo, S., Berhanu, F., Binagwa, P., Kilango, M., … Beebe, S. (2019). Genomics, genetics and breeding of common bean in Africa: A review of tropical legume project. *Plant Breeding*, *138*(4), 401–414. Online: https://doi.org/10.1111/pbr.12573

©2026, Universidad Politécnica Salesiana, Ecuador.

Myers, J. R., and Kmiecik, K. (2017). Common bean: Economic importance and relevance to biological science research. En J. R. Myers and K. Ken (Eds.), *Compendium of plant genomes* (pp. 1–20). Springer International Publishing. Online: https://doi.org/10.1007/978-3-319-63526-2_1

Nápoles, J. R. (2018). *Revisión del género Zabrotes para México (Coleoptera: Bruchidae)* (Master's thesis, Colegio de Postgraduados). Online: https://shortlink.uk/1rhxp

Nasar, S., Shaheen, H., Murtaza, G., Tinghong, T., Arfan, M., and Idrees, M. (2023). Socioeconomic evaluation of common bean (*Phaseolus vulgaris* L.) cultivation in providing sustainable livelihood to the mountain populations of Kashmir Himalayas. *Plants*, *12*(1). Online: https://doi.org/10.3390/plants12010213

Okafor, K. P., Akunne, C. E., Nwaoke, A. J., Nlemchi, C. J., Okeke, T. E., Chidi, C. A., … Offor, V. O. (2024). Evaluation of insecticidal potentials of chilli pepper (*Capsicum frutescens* L.) and guinea pepper (*Xylopia aethiopica* Dun. A. Rich.) in the control of bean weevil (*Callosobruchus maculatus* L. Fabricus). *Multidiscience: Journal of Multidisciplinary Science*, *1*(2), 51–66. Online: https://doi.org/10.59631/multidiscience.v1i2.202

Page, M. J., McKenzie, J. E., Bossuyt, P. M., Boutron, I., Hoffmann, T. C., Mulrow, C. D., … Moher, D. (2021). The PRISMA 2020 statement: An updated guideline for reporting systematic reviews. *BMJ (Clinical Research Ed.)*, *372*, n71. Online: https://doi.org/10.1136/bmj.n71

Perales-Aguilar, L., Chávez-Servia, J. L., Perales-Aguilar, L., Tafoya, F., and Perales-Segovia, C. (2020). Manejo del gorgojo de frijol almacenado, con extractos de plantas asociadas al cultivo. *Revista Mexicana de Agroecosistemas*, *7*(2), 76–84.

Permuy-Abeleira, N., Chaveco-Pérez, O., González-Ferrer, E., J. y García-Sánchez, and Hidalgo-Figueroa, N. (2008). Pérdidas de grano de frijol común en un sistema de almacenamiento tradicional. *Agricultura Técnica en México*, *34*(1), 91–100.

Pimbert, M. P., and Pierre, D. (1983). Ecophysiological aspects of bruchid reproduction. I. The influence of pod maturity and seeds of *Phaseolus vulgaris* and the influence of insemination on the reproductive activity of *Zabrotes subfasciatus*. *Ecological Entomology*, *8*(1), 87–94. Online: https://shortlink.uk/1rhxu

Ramírez Cariño, R. N., Romero Nápoles, J., Vera Graziano, J., Equihua Martínez, A., Bautista Martínez, N., Hernández Morales, J., and Aragón García, A. (2017). Demografía de brúquidos (Coleoptera: Bruchidae) asociados con *Vigna unguiculata* (L.) Walp (Fabaceae). *Acta Zoológica Mexicana*, *33*(1), 9–17.

Rámos-Archila, N. E., and Santacruz-Jiménez, S. F. (2024). *Evaluación de tres diluciones de un extracto de Allium sativum (ajo) y su posible uso como insecticida natural contra el Zabrotes subfasciatus (gorgojo)* (Bachelor's thesis). Universidad de El Salvador.

Raygoza-Martínez, A. P., Rodríguez-Herrera, S. A., Cervantes-Ortiz, F., García-Rodríguez, J. G., Rodríguez-Mercado, D., and Mendoza Elos, M. (2025). Bioinsecticidas para el control de *Zabrotes subfasciatus* Boheman y la calidad de semilla en frijol. *Revista Mexicana de Ciencias Agrícolas*, *16*(1). Online: https://doi.org/10.29312/remexca.v16i1.3086

Rees, D. P. (2018). Coleoptera. En *Integrated management of insects in stored products* (pp. 1–39). CRC Press. Online: https://doi.org/10.1201/9780203750612-1

Refaay, D. A., El-Marzoki, E. M., Abdel-Hamid, M. I., and Haroun, S. A. (2021). Effect of foliar application with *Chlorella vulgaris*, *Tetradesmus dimorphus*, and *Arthrospira platensis* as biostimulants for common bean. *Journal of Applied Phycology*, *33*(6), 3807–3815. Online: https://doi.org/10.1007/s10811-021-02584-z

Rodríguez-Sifuentes, L., Marszalek, J. E., Chuck-Hernández, C., and Serna-Saldívar, S. O. (2020). Legumes protease inhibitors as biopesticides and their defense mechanisms against biotic factors. *International Journal of Molecular Sciences*, *21*(9). Online: https://doi.org/10.3390/ijms21093322

Rodríguez Hernández, C., and López Pérez, E. (2001). Actividad insecticida e insectistática de la chilca (*Senecio salignus*) sobre *Zabrotes subfasciatus*. *Manejo Integrado de Plagas*, *62*, 45–52. Online: https://shortlink.uk/1rhxz

©2026, Universidad Politécnica Salesiana, Ecuador.

Singh, S. P. (2013). *Common bean improvement in the twenty-first century* (Vol. 7). Springer Netherlands.

Souza, S. M., Uchôa, A. F., Silva, J. R., Samuels, R. I., Oliveira, A. E. A., Oliveira, E. M., … Silva, C. P. (2010). The fate of vicilins, 7S storage globulins, in larvae and adult *Callosobruchus maculatus* (Coleoptera: Chrysomelidae: Bruchinae). *Journal of Insect Physiology*, *56*(9), 1130–1138. Online: https://doi.org/10.1016/j.jinsphys.2010.03.009

Thompson, S. N., and Suarez, R. K. (Eds.). (2009). *Encyclopedia of insects*. London, UK: Academic Press.

Tigist, S. G., Raatz, B., Assefa, A., Melis, R., Sibiya, J., Keneni, G., … Tsegaye, D. (2021). Introgression of bruchid (*Zabrotes subfasciatus*) resistance into small red common bean (*Phaseolus vulgaris*) background and validation of the BRU_00261 (snpPV0007) resistance marker. *Plant Breeding*, *140*(6), 1081–1089. Online: https://doi.org/10.1111/pbr.12969

Torres-Treminio, J. A., and Mendoza-Montoya, J. (2002). *Efecto de la fertilización mineral, orgánica y control de malezas sobre crecimiento y rendimiento en frijol común (Phaseolus vulgaris L.) en condiciones de ladera, establecido bajo dos sistemas de labranza, Ticuantepe, postrera dos mil* (Bachelor's thesis, Universidad Nacional Agraria, Managua, Nicaragua). Online: https://shortlink.uk/1wItV

Vázquez-Zamorano, R. L. (2014). *Evaluación de tres insecticidas para el control del gorgojo del maíz Sitophilus zeamays Motschulsky, bajo condiciones de laboratorio* (Bachelor's thesis, Universidad Autónoma Agraria Antonio Narro, Saltillo, Coahuila, México). Online: https://shortlink.uk/1wItZ

Wilker, J., Humphries, S., Rosas-Sotomayor, J. C., Cerna, M. G., Torkamaneh, D., Edwards, M., … Pauls, K. P. (2020). Genetic diversity, nitrogen fixation, and water use efficiency in a panel of Honduran common bean (*Phaseolus vulgaris* L.) landraces and modern genotypes. *Plants*, *9*(9). Online: https://doi.org/10.3390/plants9091238

©2026, Universidad Politécnica Salesiana, Ecuador.