



HEAVY METALS REMEDIATION WITH POTENTIAL APPLICATION IN COCOA CULTIVATION

TÉCNICAS DE REMEDIACIÓN DE METALES PESADOS CON POTENCIAL APLICACIÓN EN EL CULTIVO DE CACAO

Javier Andrés Casteblanco

Pedagogical and Technological University of Colombia UPTC, Avenida Central del Norte 39-115, 150003 Tunja, Tunja, Boyacá, Colombia.

*Author for correspondence: javierancaste@hotmail.com

Manuscript submitted on August 7, 2018. Accepted, after review on January 5, 2018. Published on March 1, 2018.

Abstract

Cacao (*Theobroma cacao*) worldwide has increased its area sown and yield per hectare, however currently producers are facing strong legislation issued by the European Union regarding the maximum levels of lead and cadmium that must have chocolates that contain an amount greater than or equal to 50% cocoa solids. Based on a review of the work carried out around the world and published in the last three years in global databases are presented, in the first instance, the problems caused in people by the consumption of food contaminated by heavy metals and the routes through which the cocoa can be contaminated, from its planting to its processing. The remediation techniques (phytoremediation and bioremediation) that have obtained good results regarding the cleaning of contaminated soils or that avoid the transfer of the contents of lead and cadmium from the soil to several Crops of commercial interest to have options of potential application in the cacao areas of Colombia or anywhere in the world. The results show the importance of implementing an integrated soil remediation system that includes the gradual incorporation of native trees, herbaceous plants, aquatic plants, biochar, bacteria and arbuscular mycorrhizae.

Keywords: Phytoremediation, bioremediation, lead, cadmium

Resumen

El cacao (*Theobroma cacao*) a nivel mundial ha aumentado su área sembrada y rendimiento por hectárea, sin embargo actualmente los productores se enfrentan a una fuerte legislación emanada por la Unión Europea respecto a los contenidos máximos de plomo y cadmio que deben tener los chocolates que contienen una cantidad mayor o igual al 50% de sólidos de cacao. En base a una revisión de los trabajos realizados alrededor del mundo y que han sido publicados en los últimos tres años en bases de datos mundiales se presentan, en primera instancia, los problemas ocasionados en las personas por el consumo de alimentos contaminados por metales pesados y las rutas a través de las cuales se puede contaminar el cacao, desde su siembra hasta su procesamiento. A continuación y dando cumplimiento a el objetivo de la revisión se muestran las técnicas de remediación (fitoremediación y bioremediación) que han obtenido buenos resultados respecto a la limpieza de suelos contaminados o que evitan la traslocación de los contenidos de plomo y cadmio del suelo a varios cultivos de interés comercial para tener opciones de potencial aplicación en las zonas cacaoteras de Colombia o cualquier parte del mundo. De los resultados obtenidos se resalta la importancia que tiene la implementación de un sistema integrado de remediación de suelos que incluya la incorporación gradual de árboles nativos, plantas herbáceas, plantas acuáticas, biochar, bacterias y micorrizas arbusculares.

Palabras claves: Fitoremediación, bioremediación, plomo, cadmio.

Suggested citation: Castebancho, J. A. 2018. Methods of remediation of heavy metals with potential application in cocoa cultivation. *La Granja: Journal of Life Sciences*. Vol. 27(1):20-33. <http://doi.org/10.17163/lgr.n27.2018.02>.

1 Introduction

Cocoa is a food product that has had a quite marked growth in producing countries due to its easy handling and the excellent economic benefits it presents (TECHNOSERVE, 2015). However, cocoa producers around the world were disturbed when the European Union announced plans to apply regulations on chocolate containing 50% or more of cocoa solids which should contain a maximum of 0.3 mg/kg of cadmium and 1 mg/kg of lead (CODEX Alimentarius, 2015).

Lead or cadmium in soils may have a natural or anthropic origin, but independently of this the plants absorb it and can accumulate it in different structures and proportions (Londño Franco, Londño Muñoz and Muñoz García, 2016; Covarrubias and Cabriales, 2017). This situation occurs in cocoa with significant accumulation in its seeds and leaves (Guerra Sierra et al., 2014) and since most of the products derived from cocoa are consumed by children it is necessary to minimize the presence of these metals in the final product (CODEX Alimentarius, 2015).

The objective of this review is to point out part of the research that has been worldwide developed over the last three years regarding the remediation techniques (phytoremediation and bioremediation) of heavy metals present in the soil, with emphasis on lead and cadmium, and that could potentially be used to manage the levels present in the producing areas of Colombia or anywhere in the world.

2 Referential Framework

The cacao tree is grown in tropical regions. It is commercially cultivated between 15 ° north and 15 ° south of the equator. The average annual temperature range goes from 23 to 30 ° C. It is cultivated from sea level to up 1,200 meters above sea level. Likewise, it needs average annual relative humidity of between 70% and 80% (Vizcaíno Cabezas and Betancourt, N.d.; Ramírez Sosa and Orrego Suaza, 2014).

World production of cocoa (*T. cacao* L.) has increased linearly from 1.2 million tons in 1961 to 4.6 million tons of dry cocoa in 2013. During the same period, the annual cocoa area had an average annual increase of 2.5%, (Vanhove, Vanhoudt and Damme, 2016) and although the world demand for

cocoa continues to increase, it also shows that production declines as a result of multiple factors, including pests and diseases, decrease in soil fertility and an increasingly hot and dry climate (TRANSMAR GROUP, 2014; Blaser et al., 2017), and this has been accompanied by an increase in international cocoa marketing prices due to the rise in oil prices, which increases the production of fertilizers, pesticides and fuels (Callejas, 2016). Even as of 2001, there was talk of a cocoa deficit that would consolidate by 2020, due to the decrease in production, generated by the poor conditions of growers in the main countries of origin, in addition to the increase in industrial consumption, which it has been stable in recent years (ProExport Colombia, 2014).

World cocoa production is absorbed mainly by three countries (60% of imports), with Switzerland being the largest importer (29.4%), followed by the United States (16.7%) and Germany with 13.9%. Most of the exports will continue to be cocoa beans, although there are attempts by the producing countries to add more value and it is to be expected that cocoa will be so scarce in 20 years that it will turn chocolate into a very expensive luxury product, the authentic chocolate will be within two decades a very expensive and scarce product that will honor its origin: *Theobroma cacao*, "the food of the gods" (Anga, 2015).

The high international prices and the demand by the processing industries has generated the global dynamics of cocoa is active, thus encouraging the expectations of the Colombian producers since the above is added the work being done by the National Federation of cacaoteros, FEDECACAO, related to the positioning and recognition of the quality of cocoa at an international level, which translates into new market opportunities for high quality grains produced by cocoa farmers from all regions of the country. In this sense, it is expected to continue increasing the presence in international markets where the Colombian cocoa is already recognized as fine flavor and aroma and it is expected to be able to obtain extra premiums on the base price, which also translates into better income and services for cocoa farmers (Cacao de Colombia, 2014).

Given this social and economic importance of cocoa, several producing and exporting countries, including Colombia, are interested in knowing the levels of heavy metals present in the cultivated areas (Jaramillo, 2013; Reyes et al., 2016; CAOBISCO/ECA/FCC, 2015) since in recent years there ha-

ve been progressive regulations of the European Community, where maximum limits are established regarding the allowable content of lead and cadmium in chocolate and other cocoa derivatives mar-

keted in Europe, Table 1, according to the regulations EC 1881/2006 and EC 488/2014 (FAO/WHO, 2014; Lanza et al., 2016; Chavez et al., 2015).

Table 1. Maximum permissible contents of lead and cadmium in chocolate (CODEX Alimentarius, 2015). The high assigned value for chocolates with % dry matter total $\geq 50\%$ stands out

Product	% of dry matter	Maximum limit of cadmium allowed**
Milk chocolate	<30%	0,10 mg/kg
Milk chocolate	$\geq 60\%$	0,30 mg/kg
chocolate	<50%	0,30 mg/kg
chocolate	$\geq 50\%$	0,80 mg/kg
Cocoa powder*	-	0,60 mg/kg

* This limit is applied to both the powder chocolate and cocoa sold to the final consumer as well as the sweetened cocoa powder used as ingredient, also sold to the final consumer (chocolate for drinking)

** These limits will take effect from January 1, 2019.

Consequently, it is essential for any country to have reference data on the content of lead and cadmium in food, especially in those that are preferably consumed by children, such as cocoa, which will allow them to establish a clear position before the international regulations that could represent a risk for the exports of the product and direct the necessary investigations to manage the incidence and severity of this problem on the health of the people (Uyttendaele, Boeck and Jacxsens, 2016), emphasizing the natural and easy techniques access for farmers known as remediation, including concepts such as phytoremediation (use of plant material that decreases the presence of metal in plants of interest) (Ahkami et al., 2017; Tahir, Yasmin and Khan, 2016; Luo et al., 2016); bioremediation (use of microorganisms alone or in combination of plant material that decreases the presence of metal in plants of interest (Brown et al., 2017; Galdames et al., 2017; Leal et al., 2017; Aggangan et al., 2017).

3 Problems associated with cadmium and lead

Recent research on contamination with lead and cadmium, and other heavy metals, in food (Al-Hossainy et al., 2017; Perryman et al., 2017; Siringkhawut et al., 2017), and its consequences on human health as conditions with lesions in preg-

nancy, gastrointestinal irritation, nausea, vomiting, kidney damage, emphysema, and lung cancer (Antoine, Fung and Grant, 2017; AbuShady et al., 2017; Shakir, Zahraw and Al-Obaidy, 2017; Amadi, Igweze and Orisakwe, 2017) clearly show the reasons about the concern that has been awakened worldwide on this subject and especially due to the progressive accumulation of these metals in the tissues of humans and animals which results in damage at the molecular genetic level (Abarshi, Dantala and Mada, 2017; Al-Gburi, Al-Tawash and Al-Lafta, 2017; Skiba, Kobylecka and Wolf, 2017; Nkansah et al., 2016). Regarding the entry of Cadmium in the body (Díaz García and Arceo, 2017) determines that it is done via gastric with a mechanism of damage that includes: dysfunction by formation of free radicals, apoptosis or activation via the caspases, protein denaturation and by decrease in transepithelial resistance. Regarding the entry of lead into the body, this same author, determines that it is via gastric, respiratory and contact, with various mechanisms that include: bone storage, mitochondrial damage and formation of free radicals, apoptosis, inflammation by activation FN-k β , RAAS and attraction of macrophages and decrease in NO production.

Chumbipuma, in 2016, describes that with respect to lead, its systemic effects are nonspecific, including adynamia, sleep disorders, headache, pain in bones and muscles, digestive symptoms (consti-

pation), abdominal pain, nausea, vomiting and decreased appetite. He also reports that acute lead toxicity occurs after respiratory exposure at high concentrations, with encephalopathy, renal failure and gastrointestinal symptoms, and finally concludes that workers exposed for a long time and without personal protection measures may present with a peripheral polyneuropathy, that predominantly affects the upper limbs, the extensor muscles that the flexors and more the dominant side, what has been called the "painter's hand" because it was presented in these workers by the use of paints with a high content of lead.

Anderson, in 2016 showed that the increase in blood lead concentrations of young university students decreases the vitality and fast linear mobility in their sperm. According to Londño Franco, Londño Muñoz and Muñoz García (2016), the World Health Organization mentions with respect to cadmium that the presentation and severity of signs, symptoms and alterations in the organism is related to the amounts, the exposure time and the metal entry route; chronic exposure shows anemia, renal dysfunction, kidney stones, osteoporosis, osteomalacia, respiratory disorders, hypertension, nervous disorders (headache, vertigo, sleep disturbance, tremors, sweating, paresis, involuntary muscle contractions), weight and appetite loss, prostate and lung cancer; in acute poisoning there is pneumonitis and pulmonary edema, gastroenteritis, nausea, vomiting, abdominal pain, diarrhea, kidney failure, and finally chromosomal aberrations, teratogenic and congenital effects and that in the kidney (renal tubules) can accumulate for up to thirty years.

Contaminants in cocoa

The metallic contaminants of cocoa can be defined as those metals, not intentionally added, that are present in cocoa as a result of production (Chavez et al., 2015; Ramtahal et al., 2016), manufacturing, elaboration, preparation (Moreno et al., 2017), treatment, packaging, transport, storage or as a product of environmental contaminations with the potential to present risks to people's health (CODEX Alimentarius, 2015; Londño Franco, Londño Muñoz and Muñoz García, 2016). These metals can be found in soils naturally or as a result of anthropogenic activity, be absorbed by plants, concentrated in seeds and taken from them by humans, which constitutes a potential risk to health (Amadi, Igweze and Orisakwe, 2017; Paul, 2017; El-Amier, Elnaggar and El-Alfy, 2017).

Various investigations have shown that during the process of cocoa benefit, there are variations in the physicochemical characteristics of the grain, depending on the type of fermenter used and the drying time, which can affect metal concentrations, as well as the quality and safety of the final product (Uyttendaele, Boeck and Jacxsens, 2016; Laires Amaiz et al., 2013).

Within this framework, it is clear that the source of contamination with lead and/ or cadmium from cocoa is diverse, the translocation from the soil to the different parts of the plant is complex and little studied and the manifestation of its presence has consequences that are palpable not only for plants but also for human beings and therefore it is necessary to update the available techniques that exist on the management of this problem in different crops, to have a frame of reference that can potentially be applied in cocoa, and thus be able to select those that are most appropriate for the environmental and natural conditions in which this crop is developed within Colombia or anywhere in the world.

4 Phytoremediation of heavy metals

The ideal plant species for phytoextraction are those that have the capacity to accumulate and tolerate high concentrations of metals in harvestable tissue (Tariq and Ashraf, 2016) and this is possible since they have the power to eliminate contaminants that persist in the environment. environment through various mechanisms such as phyllofiltration, phytostabilization, phytoextraction, phytovolatilization and phytotransformation (da Conceição Gomes et al., 2016), in addition to the containment and degradation of metals, phenolic compounds and various dyes, as well as other organic and inorganic contaminants (Tahir, Yasmin and Khan, 2016). Next, there are investigations in phytoremediation that have potential to be applied in the cultivation of cocoa, given the availability of the material with which it was worked.

? pointed out that hyperaccumulators have different accumulation potentials for different metals. The study developed by them at the laboratory level showed that corn (*Zea mays*) is a hyperaccumulator for Co and Cr after the application of a chelator such as EDTA, while sunflower (*Helianthus annuus*) proved to be a hyperaccumulator for Cd under condi-

tions similar, with a removal of 56.03%. The turnip plant (*Brassica campestris*) exhibited hyperaccumulative properties for Cr. On the other hand, the pea (*Pisum sativum*) was found to be the best accumulator of Pb without application of the EDTA chelate with a removal efficiency of 96.23% (Ojoawo, Udayakumar and Naik, 2015; Tariq and Ashraf, 2016).

It has been proven that the ability of soybean (*Glycine max*) to accumulate heavy metals present in the soil is low compared with its natural predisposition to tolerate them (Ibiang, Mitsumoto and Sakamoto, 2017), however it was evidenced that its accumulating power can be potentiated when sowing it together with material from the *Melastoma* genus (Syam et al., 2016). Another work has proven the effectiveness of soy as a cadmium hyperaccumulator, since adding up to 300 mg/Kg of nano particles of TiO₂ in the soil gives up to an additional 400% of µg of cadmium/plant (Singh and Lee, 2016).

The bark and leaves of *Moringa oleifera* can be used as an alternative absorbent for the removal of heavy metals from contaminated waters through a technique of obtaining modified citric acid (CAMOB and CAMOL) and its use in soils shows potential for research (George et al., 2016). At ground level, applications of foliar extract have been used to concentrate the lead contents of the soil where beans are grown (*Phaseolus vulgaris*) achieving a remarkable improvement in the stress caused by this element through the activation of the antioxidant system of the roots of beans (Howladar, 2014).

A study developed on the India-Pakistan border in an area cultivated with sugarcane and sorghum, where large amounts of phosphorus fertilizers are used, which are reported as sources of heavy metals in the soils, resulted in reports of the potential use of these plants as phytoextractors of lead, cadmium and copper because their Metal Bioaccumulation Factors (BAF) were around 1, which turns out to be a very good indicator (Singh and Lee, 2016). In the United States, sorghum sown in soils contaminated by heavy metals also showed its phytoextracting ability, either alone or in a mixture with arbuscular mycorrhizas of the *Azotobacter chroococcum* type (Dhawi, Datta and Ramakrishna, 2016).

The humus extracted from composted waste materials from the palm oil extraction *Elaeis guineensis*, mainly the empty bunches, improved with the addition of NPK-based fertilizers were evaluated in their fertilizing and heavy-metal phytoextrac-

tion, especially copper (Cu) in cucumber plants (*Cucurbita pepo*). The results indicate that the nutritional content of empty clusters is low, but the potassium intake is the most significant. Regarding its effect on heavy metals, it was concluded that with the addition of Nitrogen copper extraction is improved, as it had been in other works developed by Moreira et al in 2011 (Winarso, Pandutama and Purwanto, 2016).

An equally promising crop for the extraction of heavy metals is rapeseed or canola (*Brassica napus*), due to several advantages: 1. it has a great capacity to accumulate metals (2000 mg/K for cadmium), 2. It produces enough biomass, it is easy of cultivate and has great climatic adaptability and 3. It can be used in the industry.

In China, a study was carried out that showed that there are varieties that accumulate cadmium in their roots or seeds and for this reason if the variety with which hyperaccumulates in the seed should be used for the production of biodiesel and, if not, that the Hyperaccumulation occurs in the roots and must be used in the manufacture of edible vegetable oils (Fu et al., 2016). Another investigation proved that this plant has a hyperaccumulating power of zinc and that it develops very well in soils contaminated with this metal (Belouchrani et al., 2016).

In order to remove lead and cadmium from wastewater and with potential use in soils irrigated with these waters, plants of the species *Eichhornia sp.*, *Lemna sp.*, *Spirodella sp.*, *Azolla sp.*, *Pistia sp.* were used. It is concluded that the use of *Lemna sp* is promising because of its excellent hyperaccumulation of the two metals, regardless of the pH of the treated water (Verma and Suthar, 2015).

It was worked in Egypt with *Solanum nigrum* and it was proven that it has a great power of accumulation of zinc in its roots and leaves, of lead in its stems and fruits, of cobalt in its roots and fruits. It is highlighted that the levels found in their tissues were often higher than those found in the soil studied (Saad-Allah and Elhaak, 2017). It has been found that this plant is associated with arbuscular mycorrhizae of *Phylum Ascomycota* and therefore its excellent role as a cadmium hyperaccumulator (Khan et al., 2017).

In Australia, the phytoextraction produced by *Acacia pycnantha* and *Eucalyptus camaldulensis* was confirmed, concluding that *E.camaldulensis* presents in its leaves 1 500% more copper than the levels present in the soil; that the average of bioconcentration factors are higher in *E.camaldulensis* and

are very close to one (1) for copper, zinc, cadmium and lead and also the translocation factor of zinc and cadmium is higher in *Acacia pynantha* (Nirola et al., 2015). In Italy with the application during four years of an Integrated System of Phytoremediation (IPS), which includes *Acacia saligna*, *E. camaldulensis*, rhizobacteria and mycorrhizae, it was proved that phytostabilization in the soil of lead, cadmium and zinc can be produced (Guarino and Sciarrillo, 2017).

The ability of *Marsilea crenata* to accumulate lead in its roots and transfer it to its leaves and sprouts was proven by a laboratory-level experiment in tomato plants. The evaluations were made through the technique of measuring their bioelectrical responses and it was established that this plant can be used as a bioindicator of lead contamination in commercial rice or tomato crops, as its leaves turn yellow when there is excessive presence in the soil of this metal (Nurhayati, Hariadi and Lestari, 2015).

In a study developed under greenhouse conditions in Tunisia, the role of *Medicago sativa* alfalfa was evaluated and it was found that its bioconcentration factor (the relationship between the content of metals in the soil and the content in the foliage) demonstrates its potential use in any crop taking advantage of the wide range of adaptation present in this legume (Elouear et al., 2016). These results are in line with others found by (Flores-Cáceres, 2013), (Coyago and Bonilla, 2016).

5 Bioremediation of heavy metals

Bioremediation uses biological agents (microorganisms) for the complete elimination of pollutants and/or toxic substances from the environment, while the transformation of toxic pollutants into harmless forms through chemical modifications carried out by living organisms (bacteria and fungi) it is called biotransformation (Dzionic, Wojcieszynska and Guzik, 2016). In these respects, research works have been developed in recent years that have been quite successful and that show possible mechanisms of bioremediation of contaminated soils (Benyahia and Embaby, 2016), and that have the potential to be applied in the cocoa crop, given the ease of obtaining the biological material with which it was worked.

One of the difficulties that is granted to bioremediation is the long time that has to pass to see

the effects, however in works developed in different parts of the world it has been proven that the endophytic bacteria associated with hyperaccumulative plant species favor the efficiency of the bioremediation process and increase the production of vegetable biomass through three mechanisms: (1) increase in the surface of the root and the production of root hairs, (2) increase in the availability of metals, (3) increase in the transfer of soluble metals from the rhizosphere to the plant (Ahemad, 2015). Some of the species of bacteria used to improve the extraction of heavy metals and hydrocarbons are: *Burkholderia sp* (Yang et al., 2016), *Scirpus triquetus* (Chen et al., 2017), *Pseudomonas sp* J4AJ (Di Martino, 2015), *Bacillus subtilis* (Oyetibo et al., 2017), *Microbacterium sp.* SUCR140 (Soni et al., 2013), *Delftia sp.* JD2 (Ahemad, 2015).

In Brazil, a rehabilitation system was designed that included the planting of *E. camaldulensis* in furrows and *Brachiaria decumbens* between the intermediate furrows, accompanied by the sowing of arbuscular mycorrhizas of the *Glomus sp* type, which has already been used in the rehabilitation program of contaminated areas by heavy metals (?). A great power of extraction was found in this combination of materials in the *Brachiaria*, with an order of accumulation efficiency in its biomass of Zn, Cu, Cd and Pb. In addition, this work concludes that knowledge of the dynamics of these fungi in their role as rehabilitators of contaminated soils contributes to the revegetation and establishment of new plant materials in highly polluted areas (Leal et al., 2016; Nirola et al., 2015). It has also been worked with soil incorporation of arbuscular mycorrhizas *Glomus mosseae*, *Glomus intraradices*, *Glomus etunicatum* for minimization of cadmium in the soil and it was found that *Bassia indica* can be used in mixture with these fungi to decrease the dispersion of cadmium in the soil (Hashem et al., 2016).

The application of *Conocarpus biochar* in the soil at the time of sowing significantly reduced the concentrations of heavy metals in corn plants, showing a great immobilization power: 60.5% in manganese, 28% in zinc, 60% in copper, 47% copper (Al-Wabel et al., 2015).

In a laboratory test in Austria, in maize plants inoculated with the bacterium *Burkholderia phytofirmans* PsJN mixed with gravel sludge it was found that it significantly improves the fixation of heavy metals in the soil, preventing them from contaminating the vegetative material through immobiliza-

tion and stabilization processes (Touceda-González et al., 2015; Yang et al., 2016).

In India, the natural ability of native bacteria to reduce and detoxify lead, cadmium and chromium from industrial tannery effluents deposited in rivers and lands adjacent to factories was investigated. After making a biochemical characterization, it was found that *Micrococcus sp* and *Hafnia sp* have great potential for bioremediation of the aforementioned metals (Marzan et al., 2017).

The bacterium *Microbacterium oleivorans* has been studied for its ability to decompose polycyclic aromatic hydrocarbons, such as petroleum, and has also been evaluated to decontaminate land contaminated with heavy metals. Even in small dosages, its results are quite encouraging to continue testing it in different conditions and crops (Avramov et al., 2016; Xia et al., 2015).

Microorganisms have ample metabolic capacities that allow them to use different types of substrates with the objective of obtaining energy and in many cases transforming them, heavy metals are substrates that can be immobilized or transformed by these organisms using different strategies which can affect their bioavailability. This situation has allowed bioremediation techniques to be implemented that involve the use of fungi and bacteria in order to reduce the pollutant load of different environments (Beltrán-Pineda and Gómez-Rodríguez, 2016). The use of genetic engineering techniques has allowed the manipulation of microbial strains that naturally exhibit good bioremediation capacities to generate microorganisms with enhanced capacities that show promising results in in vitro and field level studies, however, the best results have been obtained when using together the capacities of plants and microorganisms in a mechanism known as symbiotic (Baghour et al., 2001). The hyperaccumulating plants (metalophitic) have the capacity to remove, reduce, transform, mineralize, degrade, volatilize or stabilize heavy metals thanks to its high capacity of accumulation in the roots and translocation different vegetative organs reaching levels of removal of up to 100% (Buta et al., 2014). Physiological and biochemical adaptations include the development of complex structures called metal-proteins or metallothioneins, which allow control in the accumulation of Cd, Cr and Hg, which also provide protection to the cell from toxic effects (Paz-Ferreiro et al., 2014). On the other hand, thanks to advances in genetic engineering many genes that

generate resistance to heavy metals have been introduced into plant cells, as is the case of the transgenic plant species that express the proteins organomercurial lyase (MerB) and MerA (mercury reductase), have a greater tolerance to organic Hg complexes and reducing Hg (II) to Hg (0). Microremediation and phytoremediation are considered promising technologies in the treatment of heavy metal contamination, their use in the field and laboratory show the biotechnological potential in the recovery of affected environments (Swain, Adhikari and Mohanty, 2013).

6 Conclusion

In nature there are numerous organisms with the ability to fix or absorb heavy metals, most of them anthropic, which are present in the soil solution and then end up contaminating the food we consume. For the cultivation of cocoa it is necessary to develop comprehensive remediation programs, including phytoremediation techniques (*B. campestris*, *M. sativa* or *E.camaldulensis*) and bioremediation (*arbuscular mycorrhizae* of the genus *Glomus*, and bacteria such as *M. oleivorans* and *B. phytofirmans*) and although the best results are achieved in relatively long times, it is necessary to start now to have the security of offering a chocolate with good flavor and aroma and that complies with the current regulations on lead and cadmium contents.

References

- Abarshi, Muawiya Musa, Edward Oyedeji Dantala and Sanusi Bello Mada. 2017. "Bioaccumulation of heavy metals in some tissues of croaker fish from oil spilled rivers of Niger Delta region, Nigeria." *Asian Pacific Journal of Tropical Biomedicine* 7(6):563–568. [online] doi: <https://doi.org/10.1016/j.apjtb.2017.05.008>.
- AbuShady, Mones M., Hanan A. Fathy, Gihan A. Fathy, Samer abd el Fatah, Alaa Ali and Mohamed A. Abbas. 2017. "Blood lead levels in a group of children: the potential risk factors and health problems." *Journal de Pediatria* 93(6):619–624. [online] doi: <https://doi.org/10.1016/j.jpmed.2016.12.006>.
- Aggangan, Nelly, Nina Cadiz, Arlene Llamado and Asuncion Raymundo. 2017. "Jatropha Curcas

- for Bioenergy and Bioremediation in Mine Tailing Area in Mogpog, Marinduque, Philippines." *Energy Procedia* 110:471–478. [online] doi: <https://doi.org/10.1016/j.egypro.2017.03.171>. 1st International Conference on Energy and Power, ICEP2016, 14–16 December 2016, RMIT University, Melbourne, Australia.
- Ahemad, Munees. 2015. "Enhancing phytoremediation of chromium-stressed soils through plant-growth-promoting bacteria." *Journal of Genetic Engineering and Biotechnology* 13(1):51–58. [online] doi: <https://doi.org/10.1016/j.jgeb.2015.02.001>.
- Ahkami, Amir H., Richard Allen White, Pubudu P. Handakumbura and Christer Jansson. 2017. "Rhizosphere engineering: Enhancing sustainable plant ecosystem productivity." *Rhizosphere* 3:233–243. [online] doi: <https://doi.org/10.1016/j.rhisph.2017.04.012>. New Understanding of Rhizosphere Processes Enabled by Advances in Molecular and Spatially Resolved Techniques.
- Al-Gburi, Hind Fadhil Abdullah, Balsam Salim Al-Tawash and Hadi Salim Al-Lafta. 2017. "Environmental assessment of Al-Hammar Marsh, Southern Iraq." *Heliyon* 3(2):e00256. [online] doi: <https://doi.org/10.1016/j.heliyon.2017.e00256>.
- Al-Hossainy, Ahmed F., Adila E. Mohamed, Fatma S.M. Hassan and M.M. Abd Allah. 2017. "Determination of cadmium and lead in perch fish samples by differential pulse anodic stripping voltammetry and furnace atomic absorption spectrometry." *Arabian Journal of Chemistry* 10:S347–S354. [online] doi: <https://doi.org/10.1016/j.arabjc.2012.09.005>.
- Al-Wabel, Mohammad I., Adel R.A. Usman, Ahmed H. El-Naggar, Anwar A. Aly, Hesham M. Ibrahim, Salem Elmaghraby and Abdulsoul Al-Omran. 2015. "Conocarpus biochar as a soil amendment for reducing heavy metal availability and uptake by maize plants." *Saudi Journal of Biological Sciences* 22(4):503–511. [online] doi: <https://doi.org/10.1016/j.sjbs.2014.12.003>.
- Amadi, Cecilia Nwadiuto, Zelinjo Nkeiruka Igweze and Orish Ebere Orisakwe. 2017. "Heavy metals in miscarriages and stillbirths in developing nations." *Middle East Fertility Society Journal* 22(2):91–100. [online] doi: <https://doi.org/10.1016/j.mefs.2017.03.003>.
- Anga, Jean-Marc. 2015. "Mercado Mundial del Cacao: Retos y oportunidades para productores de cacao en Nicaragua Managua." Organización Internacional del Cacao. . [online] available: <https://goo.gl/u1xSmD>.
- Antoine, Johann M.R., Leslie A. Hoo Fung and Charles N. Grant. 2017. "Assessment of the potential health risks associated with the aluminum, arsenic, cadmium and lead content in selected fruits and vegetables grown in Jamaica." *Toxicology Reports* 4:181–187. [online] doi: <https://doi.org/10.1016/j.toxrep.2017.03.006>.
- Avramov, Anton P., M.B. Couger, Emily L. Hartley, Craig Land, Rachel Wellendorf, Radwa A. Hanafy, Connie Budd, Donald P. French, Wouter D. Hoff and Noha Youssef. 2016. "Draft genome sequence of *Microbacterium oleivorans* strain Wellendorf implicates heterotrophic versatility and bioremediation potential." *Genomics Data* 10:54–60. [online] doi: <https://doi.org/10.1016/j.gdata.2016.09.005>.
- Baghour, Mourad, Diego A. Moreno, Gemma Villoira, Joaquín Hernández, Nicolás Castilla and Luis Romero. 2001. "Phytoextraction of Cd and Pb and Physiological Effects in Potato Plants (*Solanum Tuberosum* Var. Spunta). Importance of Root Temperature." *Journal of Agricultural and Food Chemistry* 49(11):5356–5363. [online] doi: <https://doi.org/10.1021/jf010428x>. PMID: 11714328.
- Belouchrani, Amel Souhila, Nabil Mameri, Nadia Abdi, Hocine Grib, Hakim Lounici and Nadjib Drouiche. 2016. "Phytoremediation of soil contaminated with Zn using Canola (*Brassica napus* L)." *Ecological Engineering* 95:43–49. [online] doi: <https://doi.org/10.1016/j.ecoleng.2016.06.064>.
- Beltrán-Pineda, A. E and A. M. Gómez-Rodríguez. 2016. "Biorremediación de Metales Pesados Cadmio (Cd), Cromo (Cr) y Mercurio (Hg), Mecanismos Bioquímicos e Ingeniería Genética: Una Revisión." *Revista Facultad de Ciencias Básicas* 12(2):172–197. [online] doi: <https://doi.org/10.18359/rfcb.2027>.
- Benyahia, Farid and Ahmed Shams Embaby. 2016. "Bioremediation of crude oil contaminated desert soil: Effect of biostimulation, bioaugmentation and bioavailability in biopile treatment systems." *International Journal of Environmental Re-*

- search and Public Health 13(2):1–11. [online] doi: <https://doi.org/10.3390/ijerph13020219>.
- Blaser, W.J., J. Oppong, E. Yeboah and J. Six. 2017. "Shade trees have limited benefits for soil fertility in cocoa agroforests." *Agriculture, Ecosystems & Environment* 243:83–91. [online] doi: <https://doi.org/10.1016/j.agee.2017.04.007>.
- Brown, David M., Samson Okoro, Juami van Gils, Rob van Spanning, Matthijs Bonte, Tony Hutchings, Olof Linden, Uzoamaka Egbuche, Kim Bye Bruun and Jonathan W.N. Smith. 2017. "Comparison of landfarming amendments to improve bioremediation of petroleum hydrocarbons in Niger Delta soils." *Science of The Total Environment* 596–597:284–292. [online] doi: <https://doi.org/10.1016/j.scitotenv.2017.04.072>.
- Buta, Erzsébet, Anamária Török, Bilassé Zongo, Maria Cantor, Mihai Buta, Cornelia Majdik and Árpád Csog. 2014. "Comparative Studies of the Phytoextraction Capacity of Five Aquatic Plants in Heavy Metal Contaminated Water." *Notulae Botanicae Horti Agrobotanici Cluj-Napoca* 42(1):173–179. [online] doi: <http://dx.doi.org/10.15835/nbha4219341>.
- Cacao de Colombia. 2014. "Resumen Ejecutivo Diseño de un sistema de calificación y clasificación de estándares de calidad para el cacao fino y de aroma de Colombia, p. 30." Alma café, Swiss contact. [online] available: <https://goo.gl/Z6b3b6>.
- Callejas, Edmar Salinas. 2016. "La onda larga de los precios de alimentos 1990-2016." *Economía Informa* 401:85–02. [online] doi: <https://doi.org/10.1016/j.ecin.2016.11.005>.
- CAOBISCO/ECA/FCC. 2015. *Cacao en Grano: Requisitos de calidad de la industria del chocolate y del cacao*. Editores End, M.J. y Dand, R. [online] available: <https://goo.gl/3PHLw8>.
- Chavez, E., Z.L. He, P.J. Stoffella, R.S. Mylavarapu, Y.C. Li, B. Moyano and V.C. Baligar. 2015. "Concentration of cadmium in cacao beans and its relationship with soil cadmium in southern Ecuador." *Science of The Total Environment* 533:205–214. [online] doi: <https://doi.org/10.1016/j.scitotenv.2015.06.106>.
- Chen, Xiao, Xiaoyan Liu, Xinying Zhang, Liya Cao and Xiaoxin Hu. 2017. "Phytoremediation effect of *Scirpus triqueter* inoculated plant-growth-promoting bacteria (PGPB) on different fractions of pyrene and Ni in co-contaminated soils." *Journal of Hazardous Materials* 325:319–326. [online] doi: <https://doi.org/10.1016/j.jhazmat.2016.12.009>.
- CODEX Alimentarius. 2015. "Norma general para los contaminantes y las toxinas presentes en los alimentos y piensos. CODEX STAN 193-1995." [online] available: <https://goo.gl/XqGcyo>.
- Covarrubias, Sergio and J. Cabriales. 2017. "CONTAMINACIÓN AMBIENTAL POR METALES PESADOS EN MÉXICO: PROBLEMÁTICA Y ESTRATEGIAS DE FITORREMEDIACIÓN." *Revista Internacional de Contaminación Ambiental* 33(0):7–21. [online] doi: <http://dx.doi.org/10.20937/RICA.2017.33.esp01.01>.
- Coyago, E and S. Bonilla. 2016. "Absorción de plomo de suelos altamente contaminados en especies vegetativas usadas para consumo animal y humano." *La Granja, Revista de ciencia y tecnología* 23(1):35–46. [online] doi: <https://doi.org/10.15446/acag.v66n2.58476>.
- da Conceição Gomes, Maria Angélica, Rachel Ann Hauser-Davis, Adriane Nunes de Souza and Angela Pierre Vitória. 2016. "Metal phytoremediation: General strategies, genetically modified plants and applications in metal nanoparticle contamination." *Ecotoxicology and Environmental Safety* 134:133–147. [online] doi: <https://doi.org/10.1016/j.ecoenv.2016.08.024>.
- Dhawi, Faten, Rupali Datta and Wusirika Ramakrishna. 2016. "Mycorrhiza and heavy metal resistant bacteria enhance growth, nutrient uptake and alter metabolic profile of sorghum grown in marginal soil." *Chemosphere* 157:33–41. [online] doi: <https://doi.org/10.1016/j.chemosphere.2016.04.112>.
- Di Martino, C. 2015. Estudio de bacterias del género *Pseudomonas* en la degradación de hidrocarburos y síntesis de biosurfactantes: análisis del efecto de los polihidroxialcanoatos PhD thesis Universidad de Buenos Aires, Facultad de Ciencias Exactas y Naturales. [online] available: <https://goo.gl/vDpgmg>.
- Díaz García, J. D and E. Arceo. 2017. "Daño renal asociado a metales pesados." *Revista Colombiana*

- de *Nefrología* 5(1):[online] doi: <http://dx.doi.org/10.22265/acnef.5.2.254>.
- Dzionic, Anna, Danuta Wojcieszynska and Urszula Guzik. 2016. "Natural carriers in bioremediation: A review." *Electronic Journal of Biotechnology* 23:28–36. [online] doi: <https://doi.org/10.1016/j.ejbt.2016.07.003>.
- El-Amier, Yasser A., Abdelhamid A. Elnaggar and Muhammad A. El-Alfy. 2017. "Evaluation and mapping spatial distribution of bottom sediment heavy metal contamination in Burullus Lake, Egypt." *Egyptian Journal of Basic and Applied Sciences* 4(1):55–66. [online] doi: <https://doi.org/10.1016/j.ejbas.2016.09.005>.
- FAO/WHO. 2014. "Anteproyecto de niveles máximos para el cadmio en el chocolate y productos derivados de cacao." Comisión del Codex Alimentarius. [online] available: <https://goo.gl/5MeHzF>.
- Flores-Cáceres, M. L. 2013. Caracterización de los mecanismos de respuesta antioxidante de *Medicago sativa* a metales pesados PhD thesis Universidad Autónoma de Madrid, Facultad de Ciencias. [online] available: <https://goo.gl/C54pkd>.
- Fu, D, L Jiang, A Mason, M Xiao, L Zhu, L Li, Q Zhou, C Shen and C. Huang. 2016. "Research progress and strategies for multifunctional rapeseed: A case study of China." *Journal of Integrative Agriculture* 15(8):1673–1684. [online] doi: [https://doi.org/10.1016/S2095-3119\(16\)61384-9](https://doi.org/10.1016/S2095-3119(16)61384-9).
- Galdames, A., A. Mendoza, M. Orueta, I.S. de Soto Garc a, M. S nchez, I. Virto and J.L. Vilas. 2017. "Development of new remediation technologies for contaminated soils based on the application of zero-valent iron nanoparticles and bioremediation with compost." *Resource-Efficient Technologies* 3(2):166–176. [online] doi: <https://doi.org/10.1016/j.refit.2017.03.008>.
- George, K. Shinomol, K. Bhanu Revathi, N. Deepa, C. Pooja Sheregar, T.S. Ashwini and Suchandrima Das. 2016. "A Study on the Potential of Moringa Leaf and Bark Extract in Bioremediation of Heavy Metals from Water Collected from Various Lakes in Bangalore." *Procedia Environmental Sciences* 35:869–880. [online] doi: <https://doi.org/10.1016/j.proenv.2016.07.104>. Waste Management for Resource Utilisation.
- Guarino, Carmine and Rosaria Sciarillo. 2017. "Effectiveness of in situ application of an Integrated Phytoremediation System (IPS) by adding a selected blend of rhizosphere microbes to heavily multi-contaminated soils." *Ecological Engineering* 99:70–82. [online] doi: <https://doi.org/10.1016/j.ecoleng.2016.11.051>.
- Guerra Sierra, B. E, A. X Sandoval Meza, L. S Manrique González and S. P. Barrera Rangel. 2014. "Ensayos preliminares in vitro de biosorción de cadmio cepas fúngicas nativas de suelos contaminados." *Innovaciencia Facultad de Ciencias Exactas, Físicas y Naturales* 2(1):53–58. [online] doi: <http://dx.doi.org/10.15649/2346075X.256>.
- Hashem, Abeer, Elsayed Fathi Abd Allah, Abdulaziz A. Alqarawi, Jahangir Ahmad Malik, Stephan Wirth and Dilfuza Egamberdieva. 2016. "Role of calcium in AMF-mediated alleviation of the adverse impacts of cadmium stress in *Bassia indica* [Wight] A.J. Scott." *Saudi Journal of Biological Sciences* p. [online] doi: <https://doi.org/10.1016/j.sjbs.2016.11.003>.
- Howladar, Saad M. 2014. "A novel Moringa oleifera leaf extract can mitigate the stress effects of salinity and cadmium in bean (*Phaseolus vulgaris* L.) plants." *Ecotoxicology and Environmental Safety* 100:69–75. [online] doi: <https://doi.org/10.1016/j.ecoenv.2013.11.022>.
- Ibiang, Young Basse, Hiroko Mitsumoto and Kazunori Sakamoto. 2017. "Bradyrhizobia and arbuscular mycorrhizal fungi modulate manganese, iron, phosphorus, and polyphenols in soybean (*Glycine max* (L.) Merr.) under excess zinc." *Environmental and Experimental Botany* 137:1–13. [online] doi: <https://doi.org/10.1016/j.envexpbot.2017.01.011>.
- Jaramillo, R. 2013. Metales pesados en plantaciones: Antecedentes y manejo. In *Cumbre Mundial de Cacao, Guayaquil, Ecuador 26-28 de Agosto de 2013*. [online] available: <https://goo.gl/fqvzXr>.
- Khan, Abdur Rahim, Muhammad Waqas, Ihsan Ullah, Abdul Latif Khan, Muhammad Aaqil Khan, In-Jung Lee and Jae-Ho Shin. 2017. "Culturable endophytic fungal diversity in the cadmium hyperaccumulator *Solanum nigrum* L. and their

- role in enhancing phytoremediation." *Environmental and Experimental Botany* 135:126–135. [online] doi: <https://doi.org/10.1016/j.envexpbot.2016.03.005>.
- Lanza, J. G, P. C Churión, N. J Lieno and V. H. López. 2016. "Evaluación del contenido de metales pesados en cacao (*Theobroma cacao* L.) de Santa Bárbara de Zulía, Venezuela." *SABER. Revista Multidisciplinaria del Consejo de Investigación de la Universidad de Oriente* 28:106–115. [online] available: <https://goo.gl/pu5XaQ>.
- Lares Amaiz, M, E Pérez Sira, C Álvarez Fernández, J Perozo González and S. El Khori. 2013. "Cambios de las propiedades físico-químicas y perfil de ácidos grasos en cacao de Chuao, durante el beneficio." *Agronomía Tropical* 63(1–2):37–47. [online] available: <https://goo.gl/zEao4S>.
- Leal, Aline Jaime, Edmo Montes Rodrigues, Patricia Lopes Leal, Aline Daniela Lopes Júlio, Rita de Cássia Rocha Fernandes, Arnaldo Chaer Borges and Marcos Rogério Tótola. 2017. "Changes in the microbial community during bioremediation of gasoline-contaminated soil." *Brazilian Journal of Microbiology* 48(2):342–351. [online] doi: <https://doi.org/10.1016/j.bjm.2016.10.018>.
- Leal, Patricia Lopes, Maryeimy Varón-López, Isabelle Gonçalves de Oliveira Prado, Jesse Valentim dos Santos, Claudio Roberto Fonseca Sousa Soares, Jose Oswaldo Siqueira and Fatima Maria de Souza Moreira. 2016. "Enrichment of arbuscular mycorrhizal fungi in a contaminated soil after rehabilitation." *Brazilian Journal of Microbiology* 47(4):853–862. [online] doi: <https://doi.org/10.1016/j.bjm.2016.06.001>.
- Londño Franco, L. F, P. T Londño Muñoz and F. G. Muñoz García. 2016. "Los riesgos de los metales pesados en la salud humana y animal." *Biotecnología en el Sector Agropecuario y Agroindustrial* 14:145–153. [online] available: <https://goo.gl/xRvptr>.
- Luo, Jie, Shihua Qi, X.W. Sophie Gu, Jinji Wang and Xianming Xie. 2016. "Evaluation of the phytoremediation effect and environmental risk in remediation processes under different cultivation systems." *Journal of Cleaner Production* 119:25–31. [online] doi: <https://doi.org/10.1016/j.jclepro.2016.01.043>.
- Marzan, Lolo Wal, Mehjabeen Hossain, Sohana Akter Mina, Yasmin Akter and A.M. Masudul Azad Chowdhury. 2017. "Isolation and biochemical characterization of heavy-metal resistant bacteria from tannery effluent in Chittagong city, Bangladesh: Bioremediation viewpoint." *The Egyptian Journal of Aquatic Research* 43(1):65–74. [online] doi: <https://doi.org/10.1016/j.ejar.2016.11.002>.
- Moreno, Paola Vargas, Leonardo Silva Gómez, Sandra Chaparro Acunna and Oswaldo Cárdenas. 2017. "Cadmium voltametric quantification in table chocolate produced in Chiquinquirá-Boyaca, Colombia." *Acta Agronómica* 66(2):172–177. [online] doi: <https://doi.org/10.15446/acag.v66n2.58476>.
- Nirola, Ramkrishna, Mallavarapu Megharaj, Thavamani Palanisami, Rupak Aryal, Kadiyala Venkateswarlu and Ravi Naidu. 2015. "Evaluation of metal uptake factors of native trees colonizing an abandoned copper mine a quest for phytostabilization." *Journal of Sustainable Mining* 14(3):115–123. [online] doi: <https://doi.org/10.1016/j.jsm.2015.11.001>.
- Nkansah, Marian Asantewah, Mavis Korankye, Godfred Darko and Matt Dodd. 2016. "Heavy metal content and potential health risk of geoplagic white clay from the Kumasi Metropolis in Ghana." *Toxicology Reports* 3:644–651. [online] doi: <https://doi.org/10.1016/j.toxrep.2016.08.005>.
- Nurhayati, Arry Y., Yuda C. Hariadi and P. Lestari. 2015. "Early Detection of Lead Stress on *Marsilea Crenata* Using Bioelectricity Measurement." *Procedia Environmental Sciences* 28:57–66. [online] doi: <https://doi.org/10.1016/j.proenv.2015.07.009>. The 5th Sustainable Future for Human Security (Sustain 2014).
- Ojoawo, Samson O., Gaddale Udayakumar and Pushparaj Naik. 2015. "Phytoremediation of Phosphorus and Nitrogen with *Canna x generalis* Reeds in Domestic Wastewater through NMA-MIT Constructed Wetland." *Aquatic Procedia* 4:349–356. [online] doi: <https://doi.org/10.1016/j.aqpro.2015.02.047>. INTERNATIONAL CONFERENCE ON WATER RESOURCES, COASTAL AND OCEAN ENGINEERING (ICWRCOE'15).
- Oyetibo, Ganiyu Oladunjoye, Mei-Fang Chien, Wakako Ikeda-Ohtsubo, Hitoshi Suzuki, Oluwafemi

- mi Sunday Obayori, Sunday Adekunle Adebunsoye, Matthew Olusoji Ilori, Olukayode Oladipo Amund and Ginro Endo. 2017. "Biodegradation of crude oil and phenanthrene by heavy metal resistant *Bacillus subtilis* isolated from a multi-polluted industrial wastewater creek." *International Biodeterioration & Biodegradation* 120:143–151. [online] doi: <https://doi.org/10.1016/j.aqpro.2015.02.047>.
- Paul, Dipak. 2017. "Research on heavy metal pollution of river Ganga: A review." *Annals of Agrarian Science* 15(2):278–286. [online] doi: <https://doi.org/10.1016/j.aasci.2017.04.001>.
- Paz-Ferreiro, J., H. Lu, S. Fu, A. Méndez and G. Gascó. 2014. "Use of phytoremediation and biochar to remediate heavy metal polluted soils: a review." *Solid Earth* 5(1):65–75. [online] doi: <https://doi.org/10.5194/se--5--65--2014>.
- Perryman, Shane E., Imran Lapong, Akhmad Mustafa, Rosiana Sabang and Michael A. Rimmer. 2017. "Potential of metal contamination to affect the food safety of seaweed (*Caulerpa* spp.) cultured in coastal ponds in Sulawesi, Indonesia." *Aquaculture Reports* 5:27–33. [online] doi: <https://doi.org/10.1016/j.aqrep.2016.12.002>.
- ProExport Colombia. 2014. "Oportunidades de mercado para exportar cacao colombiano." [online] available: <https://goo.gl/M52BH6>.
- Ramírez Sosa, C and S. Orrego Suaza. 2014. "Modelación económica con información espacialmente explícita de la deforestación en Urabá, Colombia, 1980-2000." *Semestre Económico* 14(29):31–51. [online] doi: <https://doi.org/10.22395/seec.v14n29a2>.
- Ramtahal, G, I Chang Yen, I Bekele, F Bekele, L Wilson, K Maharaj and L. Harryninan. 2016. "Relationships between Cadmium in Tissues of Cacao Trees and Soils in Plantations of Trinidad and Tobago." *Food and Nutrition Sciences* 7(1):37–43. [online] doi: <http://dx.doi.org/10.4236/fns.2016.71005>.
- Reyes, Yulieth, Ines Vergara, Omar Torres, Mercedes Díaz-Lagos and Edgar González-Jimenez. 2016. "Contaminación por metales pesados: Implicaciones en salud, ambiente y seguridad alimentaria." *Ingeniería Investigación y Desarrollo* 16(2):66–77. [online] doi: <http://dx.doi.org/10.1007/BF01616705>.
- Saad-Allah, Khalil M. and M.A. Elhaak. 2017. "Hyperaccumulation activity and metabolic responses of *Solanum nigrum* in two differentially polluted growth habitats." *Journal of the Saudi Society of Agricultural Sciences* 16(3):227–235. [online] doi: <https://doi.org/10.1016/j.jssas.2015.08.001>.
- Shakir, Eman, Zahraa Zahraw and Abdul Hameed M.J. Al-Obaidy. 2017. "Environmental and health risks associated with reuse of wastewater for irrigation." *Egyptian Journal of Petroleum* 26(1):95–102. [online] doi: <https://doi.org/10.1016/j.ejpe.2016.01.003>.
- Singh, Jiwan and Byeong-Kyu Lee. 2016. "Influence of nano-TiO₂ particles on the bioaccumulation of Cd in soybean plants (*Glycine max*): A possible mechanism for the removal of Cd from the contaminated soil." *Journal of Environmental Management* 170:88–96. [online] doi: <https://doi.org/10.1016/j.jenvman.2016.01.015>.
- Siriangkhwut, Watsaka, Patcharee Sittichan, Kraingkrai Ponghong and Piyanete Chantiratikul. 2017. "Quality assessment of trace Cd and Pb contaminants in Thai herbal medicines using ultrasound-assisted digestion prior to flame atomic absorption spectrometry." *Journal of Food and Drug Analysis* 25(4):960–967. [online] doi: <https://doi.org/10.1016/j.jfda.2016.12.011>.
- Skiba, E, J Kobilecka and W. Wolf. 2017. "Influence of 2,4-D and MCPA herbicides on uptake and translocation of heavy metals in wheat (*Triticum aestivum* L.)." *Environmental Pollution* 220:882–890. [online] doi: <https://doi.org/10.1016/j.envpol.2016.10.072>.
- Soni, S. K, R Singh, A Awasthi and A. Kalra. 2013. "Cr(VI)-reducing *Microbacterium* sp. strain SUCR140 enhances growth and yield of *Zea mays* in Cr(VI) amended soil through reduced chromium toxicity and improves colonization of arbuscular mycorrhizal fungi." *Environmental Science and Pollution Research* 21(3):1971–1979. [online] doi: <https://doi.org/10.1007/s11356--013--2098--7>.
- Swain, G, S Adhikari and P. Mohanty. 2013. "Phytoremediation of Copper and Cadmium from

- Water Using Water Hyacinth, *Eichhornia crassipes*." *International Journal of Agricultural Science and Technology* 2(1):[online] doi: <https://doi.org/10.14355/ijast.2014.0301.01>.
- Syam, Netty, Tatik Wardiyati, Muhammad Dawam Maghfoer, Eko Handayanto, Bahtiar Ibrahim and Aminah Muchdar. 2016. "Effect of Accumulator Plants on Growth and Nickel Accumulation of Soybean on Metal-contaminated Soil." *Agriculture and Agricultural Science Procedia* 9:13–19. [online] doi: <https://doi.org/10.1016/j.aaspro.2016.02.109>. International Conference on Food, Agriculture and Natural Resources, IC-FANRes 2015.
- Tahir, Uruj, Azra Yasmin and Umair Hassan Khan. 2016. "Phytoremediation: Potential flora for synthetic dyestuff metabolism." *Journal of King Saud University - Science* 28(2):119–130. [online] doi: <https://doi.org/10.1016/j.jksus.2015.05.009>.
- Tariq, Saadia R. and Azka Ashraf. 2016. "Comparative evaluation of phytoremediation of metal contaminated soil of firing range by four different plant species." *Arabian Journal of Chemistry* 9(6):806–814. [online] doi: <https://doi.org/10.1016/j.arabjc.2013.09.024>.
- TECHNOSERVE. 2015. *Construcción de un modelo de agricultura competitiva en Colombia*. [online] available: <https://goo.gl/sRXTEm>.
- Touceda-González, María, Günter Brader, Livio Antonielli, Vivek Balakrishnan Ravindran, Georg Waldner, Wolfgang Friesl-Hanl, Erika Corretto, Andrea Campisano, Michael Pancher and Angela Sessitsch. 2015. "Combined amendment of immobilizers and the plant growth-promoting strain *Burkholderia phytofirmans* PsJN favours plant growth and reduces heavy metal uptake." *Soil Biology and Biochemistry* 91:140–150. [online] doi: <https://doi.org/10.1016/j.agee.2017.04.007>.
- TRANSMAR GROUP. 2014. *Situación y perspectivas del mercado mundial de cacao*. World Bank Webinar. [online] available: <https://goo.gl/EvhcY>.
- Uyttendaele, Mieke, Elie De Boeck and Liesbeth Jacxsens. 2016. "Challenges in Food Safety as Part of Food Security: Lessons Learnt on Food Safety in a Globalized World." *Procedia Food Science* 6:16–22. [online] doi: <https://doi.org/10.1016/j.profoo.2016.02.003>. International Conference of Sabaragamuwa University of Sri Lanka 2015 (IC-SUSL 2015).
- Vanhove, Wouter, Niels Vanhoudt and Patrick Van Damme. 2016. "Effect of shade tree planting and soil management on rehabilitation success of a 22-year-old degraded cocoa (*Theobroma cacao* L.) plantation." *Agriculture, Ecosystems & Environment* 219:14–25. [online] doi: <https://doi.org/10.1016/j.agee.2015.12.005>.
- Verma, Rashmi and Surindra Suthar. 2015. "Lead and cadmium removal from water using duckweed *Lemna gibba* L.: Impact of pH and initial metal load." *Alexandria Engineering Journal* 54(4):1297–1304. [online] doi: <https://doi.org/10.1016/j.aej.2015.09.014>.
- Vizcaíno Cabezas, D. A and R. Betancourt. N.d. *Manual de Aplicabilidad de Buenas Prácticas Agrícolas para Cacao*. Agrocalidad, Ecuador. [online] available: <https://goo.gl/XsmQy5>.
- Winarso, Sugeng, Martinus H. Pandutama and Lutfi Dwi Purwanto. 2016. "Effectivity of Humic Substance Extracted from Palm Oil Compost as Liquid Fertilizer and Heavy Metal Bioremediation." *Agriculture and Agricultural Science Procedia* 9:146–157. [online] doi: <https://doi.org/10.1016/j.aaspro.2016.02.110>. International Conference on Food, Agriculture and Natural Resources, IC-FANRes 2015.
- Xia, Ye, Seth DeBolt, Jamin Dreyer, Delia Scott and Mark A. Williams. 2015. "Characterization of culturable bacterial endophytes and their capacity to promote plant growth from plants grown using organic or conventional practices." *Frontiers in Plant Science* 6:490. [online] doi: <https://doi.org/10.3389/fpls.2015.00490>.
- Yang, Zhihui, Zhi Zhang, Liyuan Chai, Yong Wang, Yi Liu and Ruiyang Xiao. 2016. "Bioleaching remediation of heavy metal-contaminated soils using *Burkholderia* sp. Z-90." *Journal of Hazardous Materials* 301:145–152. [online] doi: <https://doi.org/10.1016/j.jhazmat.2015.08.047>.